

# General Mathematics and CPS II

## Exercise 18

April 11, 2014

1. Use the simplex method on the following linear programming problem, applying the method from class to find the initial set of basic variables.

Minimize

$$z = -x_1 - x_2 - x_3$$

subject to

$$\begin{aligned}x_1 + 2x_2 + x_3 &= 4, \\2x_1 + x_2 + 2x_3 &= 10, \\ \mathbf{x} &\geq 0.\end{aligned}$$

2. The *primal* form of a linear programming problem is

$$\begin{aligned}\text{minimize } & \mathbf{c}^T \mathbf{x} \\ \text{subject to } & A\mathbf{x} = \mathbf{b}, \mathbf{x} \geq 0.\end{aligned}\tag{P}$$

The corresponding dual problem reads

$$\begin{aligned}\text{maximize } & \mathbf{y}^T \mathbf{b} \\ \text{subject to } & \mathbf{y}^T A \leq \mathbf{c}^T.\end{aligned}\tag{D}$$

Here,  $A$  is an  $m \times n$  matrix,  $\mathbf{x}, \mathbf{c} \in \mathbb{R}^n$ , and  $\mathbf{y}, \mathbf{b} \in \mathbb{R}^m$ .

Show that if  $\mathbf{x}$  solves (P) and  $\mathbf{y}$  solves (D), then

$$\mathbf{y}^T \mathbf{b} \leq \mathbf{c}^T \mathbf{x}.$$

Conclude that the primal problem does not have a finite minimum if and only if the feasible region of the dual problem is empty.

3. In the notation of the previous question, show that if  $\mathbf{x}$  is feasible for problem (P) and  $\mathbf{y}$  is feasible for problem (D), and if furthermore

$$\mathbf{y}^T \mathbf{b} = \mathbf{c}^T \mathbf{x},$$

then  $\mathbf{x}$  solves (P) and  $\mathbf{y}$  solves (D).