Applied Differential Equations and Modeling

Homework 2

Due in class Tuesday, February 19, 2019

1. Consider the differential equation

$$y' = (1 - 2x)/y$$
, $y(1) = -2$.

Solve this initial value problem, plot the graph of the solution, and state on which interval the solution is defined.

2. Consider the differential equation

$$\frac{\mathrm{d}y}{\mathrm{d}x} = \frac{y - 4x}{x - y}$$

(a) Show that, setting v = x/y, that this differential equation can be written in the form

$$v + x \frac{\mathrm{d}v}{\mathrm{d}x} = \frac{v - 4}{1 - v} \,.$$

- (b) Show that this new form of the equation is separable, and solve the equation. Write, as your final answer, y as a function of x.
- (c) Draw the direction field in the x-y-plane, and a few integral curves (i.e., solution curves to the differential equation which you found in part b). Remark: Note that the computation above showed that the direction field depends only on y/x; this should be clearly seen in your answer. Such equations are sometimes called homogeneous.
- 3. Suppose that a tank containing a certain liquid has an outlet near the bottom. Let h(t) be the height of the liquid surface above the outlet at time t. Torricelli's principle states that the outflow velocity v at the outlet is equal to the velocity of a particle falling freely (with no drag) from height h.
 - (a) Show that $v = \sqrt{2gh}$, where g is the constant of gravitational acceleration.
 - (b) By equating the rate of outflow to the rate of change of volume in the tank, show that h(t) satisfies the differential equation

$$A(h) \frac{\mathrm{d}h}{\mathrm{d}t} = -\alpha \, a \sqrt{2gh} \,,$$

where A(h) is the area of the cross section of the tank at height h and a is the area of the outlet. The constant α is a contraction coefficient that accounts for the observed fact that the cross section of the (smooth) outflow stream is smaller that a. The value of α for water is about 0.6.

- (c) Consider a water tank in the form of a cylinder with circular cross section. Its height is 3 m above the outlet, its radius is 1 m, and the radius of the circular outlet is $\frac{1}{10}$ m. If the tank is initially full of water, determine how long it takes to drain the tank down to the level of the outlet.
- 4. A ball with mass 0.15 kg is thrown upward with initial velocity 20 m/s from a position 30 m above ground. The ball is subject to gravitational force and a force due to to air resistance of magnitude k |v| with k = 1/30 kg/s.
 - (a) Find the maximum height above the ground that the ball reaches.
 - (b) Find the time that the ball hits the ground.
 - (c) Plot the graphs of velocity and position versus time.
- 5. For each of the following initial value problems, find the solution and state how the interval of existence of the solution depends on the initial value a.
 - (a) $y' = 2ty^2$, y(0) = a
 - (b) $y' = -4\frac{t}{y}$, y(0) = a
 - (c) $y' + y^3 = 0$, y(0) = a