

Finite rank vector bundles on inductive limits of grassmannians^{*†}

Joseph Donin and Ivan Penkov

November 3, 2006

To the memory of Andrey Tyurin

Abstract

If \mathbb{P}^∞ is the projective ind-space, i.e. \mathbb{P}^∞ is the inductive limit of linear embeddings of complex projective spaces, the Barth-Van de Ven-Tyurin (BVT) Theorem claims that every finite rank vector bundle on \mathbb{P}^∞ is isomorphic to a direct sum of line bundles. We extend this theorem to general sequences of morphisms between projective spaces by proving that, if there are infinitely many morphisms of degree higher than one, every vector bundle of finite rank on the inductive limit is trivial. We then establish a relative version of these results, and apply it to the study of vector bundles on inductive limits of grassmannians. In particular we show that the BVT Theorem extends to the ind-grassmannian of subspaces commensurable with a fixed infinite dimensional and infinite codimensional subspace in \mathbb{C}^∞ . We also show that, for a class of twisted ind-grassmannians, every finite rank vector bundle is trivial.

2000 AMS Subject Classification: Primary 32L05, 14J60, Secondary 14M15.

1 Introduction

About 30 years ago the study of "infinitely extendable" vector bundles of finite rank on projective spaces and grassmannians was initiated. In particular the following remarkable theorem was proved: any finite rank vector bundle on the infinite complex projective space \mathbb{P}^∞ (or equivalently, any finite rank vector bundle on \mathbb{P}^n which admits an extension to \mathbb{P}^m for large enough $m > n$) is isomorphic to a direct sum of line bundles. For rank two bundles this was established by W. Barth and A. Van de Ven in [BV], and for arbitrary finite rank bundles the theorem was proved by A. Tyurin in [T]. In what follows we refer to this result as to the Barth-Van de Ven-Tyurin Theorem, or as to the BVT Theorem. Some first steps were made also towards understanding finite rank vector bundles on infinite

*Work supported in part by Israel Academy of Sciences Grant no. 8007/99-03 and the Max Planck Institute for Mathematics, Bonn

†Work supported in part by an NSF grant and the Max Planck Institute for Mathematics, Bonn

grassmannians: R. Hartshorne conjectured that every finite rank bundle on an infinite grassmannian $G(k, \infty)$ is homogeneous, see [BV]. For rank two bundles this conjecture is proved in [BV], and in the general case the conjecture is proved by E. Sato, [S2]. Sato established also a partial analog of Hartshorne's conjecture for the infinite grassmannians of the classical simple groups, and reproved the BVT Theorem, see [S2] and [S1].

The purpose of the present note is to revive this discussion and relate it with the more recent discussion of homogeneous ind-spaces of locally linear ind-groups, see [DPW], [DP] and the references therein. Our starting point is an infinite sequence

$$G_1 \subset G_2 \subset \dots$$

of complex linear algebraic groups and a subsequence

$$P_1 \subset P_2 \subset \dots$$

of parabolic subgroups. This yields a sequence of morphisms

$$G_1/P_1 \longrightarrow G_2/P_2 \longrightarrow \dots \tag{1}$$

Let G/P denote the inductive limit of (1). We restrict ourselves here to the case when $G_N \simeq GL(n_N)$ and P_N are maximal parabolic subgroups, i.e. G_N/P_N are grassmannians.

The study of line bundles on G/P gives some first hints on what general finite rank bundles on G/P might look like. An essential difference with the cases studied in [BV], [T], [S1], and [S2] is that the restriction maps

$$\text{Pic}(G_N/P_N) \longrightarrow \text{Pic}(G_{N-1}/P_{N-1})$$

on the Picard groups induced by inclusions in (1) are only injective and in general not surjective. Therefore $\text{Pic}(G/P)$, the Picard group of the inductive limit, is isomorphic to \mathbb{Z} or equals zero. In the first case we call the sequence (1) *linear* and in the second case we call it *twisted*.

Here is a brief description of our results. The simplest case for a sequence (1) is when all G_N/P_N are projective spaces. Consider more generally an arbitrary sequence of morphisms of projective spaces

$$\mathbb{P}^{i_1} \xrightarrow{\varphi_1} \mathbb{P}^{i_2} \xrightarrow{\varphi_2} \dots \tag{2}$$

If (2) is linear, i.e. all but finitely many Picard groups map isomorphically, the inductive limit is isomorphic to the projective ind-space \mathbb{P}^∞ . Here the BVT Theorem claims that any finite rank bundle is isomorphic to a direct sum of line bundles. If (2) is twisted, i.e. the Picard group of the inductive limit \mathbb{P}_{tw}^∞ of (2) equals zero, the problem of describing all finite rank vector bundles on \mathbb{P}_{tw}^∞ was not posed in the 70's.

Our first main result (Theorem 3.1) claims that every finite rank vector bundle on \mathbb{P}_{tw}^∞ is trivial. This theorem, together with the BVT Theorem, can be extended to the relative case and yields a complete description of finite rank vector bundles on any inductive limit of relative projective spaces.

In Section 4 we apply the above result to the study of finite rank vector bundles on inductive limits of grassmannians $G(k_N, n_N)$. We consider two types of morphisms of grassmannians in (1): standard inclusions and a certain class of twisted homogeneous morphisms which we call twisted extensions. For standard inclusions we show that, if $\lim_{N \rightarrow \infty} k_N = \lim_{N \rightarrow \infty} (n_N - k_N) = \infty$, any finite rank vector bundle on the inductive limit is isomorphic to a direct sum of line bundles. An interesting ind-variety which arises as the inductive limit of a sequence of standard inclusions satisfying the above condition is the ind-grassmanian $G(V, \infty)$ of subspaces $V' \subset \mathbb{C}^\infty$ commensurable with a fixed infinite dimensional and infinite codimensional subspace $V \subset \mathbb{C}^\infty$. Therefore, any finite rank vector bundle on $G(V, \infty)$ is isomorphic to a direct sum of line bundles. Finally, we prove that in the case of twisted extensions of grassmannians every finite rank vector bundle on the inductive limit is trivial.

Acknowledgements. We are grateful to A. Tyurin for a helpful discussion on the topic of this paper several weeks before he suddenly passed away in October 2002. We thank also Z. Ran for his comments and for making us aware of E. Sato's work, and the referee for helpful comments.

2 Preliminary results

2.1 Notation

The ground field is \mathbb{C} and we work in the category of complex analytic spaces. A *vector bundle* always means a vector bundle of finite rank. If E is a vector bundle, we denote by E^* the dual vector bundle. For the tensor power of a line bundle E we write simply E^k . More generally, the convention $E^{-k} = (E^*)^k$ enables us to assume that $k \in \mathbb{Z}$. By $H^i(E)$ we denote the i -th cohomology group of the sheaf of local sections of E , and put $h^i(E) = \dim H^i(E)$, and $\chi(E) := \sum_i (-1)^i h^i(E)$.

By $G(k, n)$ we denote the grassmannian of k -dimensional subspaces in \mathbb{C}^n . For $k = 1$ we have the projective space $\mathbb{P}^{n-1} := G(1, n)$. The grassmannian $G(k, n)$ is isomorphic to $G(n - k, n)$ via the isomorphism

$$\{V \subset \mathbb{C}^n\} \mapsto \{V^\perp \subset \mathbb{C}^n\},$$

where \mathbb{C}^n is identified with its dual space. Under a *projective subspace* of $G(k, n)$ we understand the set of k -dimensional subspaces $V \subset \mathbb{C}^n$ such that $U \subset V \subset W$, where $U \subset W$ are fixed subspaces of \mathbb{C}^n with $\dim U = k - 1$, $\dim W > k$, or $\dim W = k + 1$, $\dim U < k$. The projective subspace is a *line* if $\dim W - \dim V = 2$.

Let S_k be the vector bundle on $G(k, n)$ with fiber V at the point $V \in G(k, n)$. There is a canonical inclusion $S_k \subset \tilde{\mathbb{C}}^n$, where $\tilde{\mathbb{C}}^n$ denotes the trivial bundle on $G(k, n)$ with fiber \mathbb{C}^n . We set $S_{n-k} := (\tilde{\mathbb{C}}^n/S_k)^*$. By definition, S_k and S_{n-k} are the *tautological*

bundles on $G(k, n)$. The Picard group $\text{Pic}(G(k, n))$ is isomorphic to \mathbb{Z} , and both maximal exterior powers $\wedge^k S_k$ and $\wedge^{n-k} S_{n-k}$ are isomorphic generators of $\text{Pic}(G(k, n))$. We set $\mathcal{O}_{G(k, n)}(-1) := \wedge^k S_k \simeq \wedge^{n-k} S_{n-k}$ and $\mathcal{O}_{G(k, n)}(m) := \mathcal{O}_{G(k, n)}(-1)^{-m}$ for $m \in \mathbb{Z}$.

An *ind-space* is the union $\cup_N X_N$ of analytic spaces X_N related by closed immersions

$$X_1 \xrightarrow{\varphi_1} X_2 \xrightarrow{\varphi_2} \dots \quad (3)$$

For instance, the projective ind-space \mathbb{P}^∞ is the union $\cup_N \mathbb{P}^N$ where $\mathbb{P}^N \subset \mathbb{P}^{N+1}$ is the standard closed immersion. A *morphism of ind-spaces* $\varphi : X \rightarrow Y$ is a map whose restriction to each X_N is a morphism of X_N into $Y_{j(N)}$ for some $j(N)$. In this paper we will more generally consider sequences (3) of arbitrary morphisms. A *vector bundle* on the system (3) is a collection $\{E_N\}$ of vector bundles E_N on X_N such that $\varphi_N^* E_{N+1} = E_N$. If (3) determines an ind-space X , we speak of a vector bundle on X .

2.2 The Barth-Van de Ven-Tyurin Theorem

Theorem 2.1. *If $X \simeq \mathbb{P}^1$ or $X \simeq \mathbb{P}^\infty$, every vector bundle E on X is isomorphic to a unique direct sum of line bundles, i.e. $E \simeq \bigoplus_j \mathcal{O}_X(d_j)$ for some unique integers $d_1 \geq \dots \geq d_{\text{rk } E}$.*

For $X \simeq \mathbb{P}^1$ this is a classical result due to A. Grothendieck. For $X \simeq \mathbb{P}^\infty$ the theorem has been proved by A. Tyurin in [T] (and earlier by W. Barth and A. Van de Ven, [BV], for vector bundles of rank two). E. Sato also presents a proof in [S1].

We will make extensive use of Theorem 2.1. In particular, if $l \subset Y$ is a smooth rational curve in a complex manifold or ind-space Y and E is a vector bundle on Y , we call the numbers $d_1 \geq \dots \geq d_{\text{rk } E}$, together with the multiplicities with which they occur in the isomorphism $E|_l \simeq \bigoplus_j \mathcal{O}_l(d_j)$, the *splitting data* of $E|_l$. We set $D := d_1 - d_{\text{rk } E}$. When we need to emphasize the dependence of d_j and D on l , we write $d_j(E|_l)$ and $D(E|_l)$.

We will use the same convention when Y is an ind-space and l is replaced by a smooth ind-subspace $P \subset Y$ isomorphic to \mathbb{P}^∞ . Note that, if E' is a vector bundle on $X \simeq \mathbb{P}^1$ or $X \simeq \mathbb{P}^\infty$ with splitting data $d_1 \geq \dots \geq d_{\text{rk } E}$, and ν_1 is the multiplicity of d_1 , there is a unique subbundle E'_1 of E' isomorphic to the direct sum of ν_1 copies of $\mathcal{O}_X(d_1)$. Indeed, consider the twisted bundle $E' \otimes \mathcal{O}_X(-d_1)$ and its unique maximal subbundle E'' generated by global sections. Then $E'_1 = E'' \otimes \mathcal{O}_X(d_1)$.

We say that a vector bundle E on a grassmannian $G(k, n)$ is *uniform*, if the splitting data $d_1(E|_l), \dots, d_{\text{rk } E}(E|_l)$ for a line $l \subset G(k, n)$ does not depend on the choice of l . The bundle E is *linearly trivial* if its restriction $E|_l$ is trivial for any line $l \subset G(k, n)$. A linearly trivial bundle is necessarily trivial: for $k = 1$ this is a well-known result, and an induction argument on k yields the result also for $G(k, n)$, see for instance [P].

2.3 Local rigidity of direct sums of line bundles on \mathbb{P}^N

Let \mathcal{S} be an analytic space and let π_1 and π_2 denote respectively the projections of $\mathbb{P}^1 \times \mathcal{S}$ onto the first and the second factor. If E is a vector bundle on $\mathbb{P}^1 \times \mathcal{S}$, $E(x)$ denotes the restriction of E to $\pi_2^{-1}(x) \simeq \mathbb{P}^1$. In [Do1] the following statement is proved.

Proposition 2.2. *Let E be a vector bundle on $\mathbb{P}^1 \times \mathcal{S}$, where \mathcal{S} is a connected analytic space. Fix $x_0 \in \mathcal{S}$. Then*

a) *there exists an open neighborhood $U \ni x_0$ such that $d_1(E(x)) \leq d_1(E(x_0))$ for $x \in U$, i.e. $d_1(E(x))$ is an upper semicontinuous function on \mathcal{S} ;*

b) *$\sum_{i=1}^{\text{rk } E(x)} \nu_i(E(x)) d_i(E(x)) = c_1(E(x))$, where $c_1(E(x))$ is the first Chern class of $E(x)$ and $\nu_i(E(x))$ is the multiplicity of $d_i(E(x))$ (in particular $\sum_{i=1}^{\text{rk } E(x)} \nu_i(E(x)) d_i(E(x))$ does not depend on $x \in \mathcal{S}$).*

Proposition 2.2, applied to E^* implies that $d_{\text{rk } E}(E(x)) = -d_1(E^*(x))$ is a lower semicontinuous function on \mathcal{S} , hence $D(E(x)) = d_1 - d_{\text{rk } E}$ is also an upper semicontinuous function on \mathcal{S} .

Proposition 2.3. *Let E be a vector bundle on \mathbb{P}^N , $N > 1$, which is isomorphic to a direct sum of line bundles. Then E is locally rigid.*

Proof. Note that $H^1(\text{End}(E)) = 0$. This follows from the fact that $\text{End}(E)$ is isomorphic to a direct sum of line bundles, as $H^1(\mathcal{L}) = 0$ for any line bundle \mathcal{L} on \mathbb{P}^N for $N > 1$. Therefore the parameter space of the versal deformation of E consists of a single reduced point, see [Do2]. \square

2.4 Estimates related to vector bundles on $\mathbb{P}^1 \times \mathbb{P}^1$

In the case of $\mathbb{P}^1 \times \mathbb{P}^1$ we call the fibers of π_1 (respectively, π_2) vertical (resp., horizontal) sections of $\mathbb{P}^1 \times \mathbb{P}^1$, and for any vector bundle E on $\mathbb{P}^1 \times \mathbb{P}^1$ we denote the twisted bundle $E \otimes \pi_1^*(\mathcal{O}(k)) \otimes \pi_2^*(\mathcal{O}(l))$ by $E(k, l)$.

Lemma 2.4. *Let E be a vector bundle on $\mathbb{P}^1 \times \mathbb{P}^1$. Let P be a vertical section and S be a horizontal section of $\mathbb{P}^1 \times \mathbb{P}^1$. Then $d_1(E|_P), d_1(E|_S) \geq -1$ implies*

$$h^0(E) \leq (\text{rk } E)(d_1(E|_P) + 1)(d_1(E|_S) + 1). \quad (4)$$

If $d_1(E|_P) < 0$ or $d_1(E|_S) < 0$, we have $h^0(E) = 0$.

Proof. The second statement follows immediately from Proposition 2.2, since $h^0(F) = 0$ for any vector bundle F on \mathbb{P}^1 when $d_1(F) < 0$. If $d_1(E|_P)$ or $d_1(E|_S)$ equals -1 , then (4) is obvious, therefore we can assume that $d_1(E|_P), d_1(E|_S) \geq 0$.

For any k there is the exact sequence

$$0 \longrightarrow E(0, -k-1) \longrightarrow E(0, -k) \longrightarrow E|_S \longrightarrow 0.$$

The corresponding cohomology sequence gives

$$h^0(E(0, -k)) - h^0(E(0, -k-1)) \leq h^0(E|_S) \leq (\text{rk } E)(d_1(E|_S) + 1).$$

Summation from k to zero yields

$$h^0(E) - h^0(E(0, -k-1)) \leq (\text{rk } E)(k+1)(d_1(E|_S) + 1). \quad (5)$$

When $k = d_1(E|_P)$ we have, $h^0(E(0, -k-1)) = 0$, therefore (5) implies (4). \square

Lemma 2.5. *Let E , P , and S be as in Lemma 2.4. Suppose $d_1(E|_P), d_1(E|_S) \geq 0$ and $d_{\text{rk } E}(E|_P), d_{\text{rk } E}(E|_S) \leq 0$. If P' is another vertical section of $\mathbb{P}^1 \times \mathbb{P}^1$, then*

$$D(E|_{P'}) \leq 4(\text{rk } E)(D(E|_P) + 2)(D(E|_S) + 1) - \chi(E) - \chi(E^*). \quad (6)$$

Proof. From the cohomology sequence of the exact sequence

$$0 \longrightarrow E \longrightarrow E(1, 0) \longrightarrow E|_{P'} \longrightarrow 0$$

we have

$$h^0(E) + h^0(E|_{P'}) \leq h^0(E(1, 0)) + h^1(E).$$

This, together with the equality $h^2(E) = h^0(E^*(-2, -2))$ (Serre duality), gives

$$d_1(E|_{P'}) \leq h^0(E|_{P'}) \leq h^0(E(1, 0)) - \chi(E) + h^0(E^*(-2, -2)). \quad (7)$$

Note that $d_1((E^*(-2, -2)|_P)) = -d_{\text{rk } E}(E|_P) - 1$ and $d_1((E^*(-2, -2)|_S)) = -d_{\text{rk } E}(E|_S) - 1$, so Lemma 2.4 yields

$$h^0(E^*(-2, -2)) \leq (\text{rk } E)(-d_{\text{rk } E}(E|_P))(-d_{\text{rk } E}(E|_S)). \quad (8)$$

Combining (7) and (8) we have

$$d_1(E|_{P'}) \leq (\text{rk } E)(d_1(E|_P) + 2)(d_1(E|_S) + 1) - \chi(E) + (\text{rk } E)(-d_{\text{rk } E}(E|_P))(-d_{\text{rk } E}(E|_S)).$$

Since $d_1(E|_P), -d_{\text{rk } E}(E|_P) \leq D(E|_P)$ and $d_1(E|_S), -d_{\text{rk } E}(E|_S) \leq D(E|_S)$, we obtain

$$d_1(E|_{P'}) \leq 2(\text{rk } E)(D(E|_P) + 2)(D(E|_S) + 1) - \chi(E). \quad (9)$$

By adding to (9) the analogous inequality for E^* we obtain (6). \square

3 The case of a twisted projective ind-space

Our first main result is the following theorem.

Theorem 3.1. *Let*

$$\mathbb{P}^{i_1} \xrightarrow{\varphi_1} \mathbb{P}^{i_2} \xrightarrow{\varphi_2} \dots$$

be a twisted sequence of projective spaces, i.e. $\deg \varphi_N > 1$ for infinitely many N , and let $E = \{E_N\}$ be a vector bundle on this inductive system. Then E is trivial, i.e. all E_N are trivial.

Lemma 3.2. *Let E be as in Theorem 3.1. Then all Chern classes $c_q(E_N)$ equal zero.*

Proof. By definition, $c_q(E_N) \in H^{2q}(\mathbb{P}^{i_N}, \mathbb{Z})$. For each $j > N$, the homomorphism

$$H^{2q}(\mathbb{P}^{i_j}, \mathbb{Z}) \longrightarrow H^{2q}(\mathbb{P}^{i_N}, \mathbb{Z}) \quad (10)$$

induced by the composition $\varphi_{j,N} := \varphi_N \circ \dots \circ \varphi_j$ maps $c_q(E_j)$ to $c_q(E_N)$. On the other hand, we can identify $H^{2q}(\mathbb{P}^{i_N}, \mathbb{Z})$ and $H^{2q}(\mathbb{P}^{i_j}, \mathbb{Z})$ in a standard way with \mathbb{Z} , and (10) considered as an endomorphism of \mathbb{Z} is nothing but multiplication by $(\deg \varphi_j) \cdot \dots \cdot (\deg \varphi_N)$. Therefore $c_q(E_N) \in \mathbb{Z}$ is divisible by $(\deg \varphi_j) \cdot \dots \cdot (\deg \varphi_N)$ for all $j > N$. According to our assumption, $\lim_{j \rightarrow \infty} ((\deg \varphi_j) \cdot \dots \cdot (\deg \varphi_N)) = \infty$, consequently, $c_q(E_N) = 0$. \square

Proof of Theorem 3.1.

Without restriction of generality we assume that $i_1 = 1$. Fix N with $i_N > 2$ and set

$$D_N := \max_{l'} D(E_N|_{l'}),$$

where l' runs over all lines in \mathbb{P}^{i_N} . Let l be a line in \mathbb{P}^{i_N} such that $D(E|_l) = D_N$, and Q be a projective subspace in \mathbb{P}^{i_N} of dimension $i_N - 2$ not intersecting l . We can choose homogeneous coordinates z_0, \dots, z_{i_N} in \mathbb{P}^{i_N} such that l is defined by $z_i = 0, i \geq 2$, and Q is defined by $z_0 = z_1 = 0$. Furthermore, we fix a morphism $f : \mathbb{P}^1 \rightarrow \mathbb{P}^{i_N}$ of degree $\deg \varphi_{1,n}$ such that

$$f(\mathbb{P}^1) \cap l = \emptyset, \quad f(\mathbb{P}^1) \cap Q = \emptyset, \quad (11)$$

and

$$D(f^*E_N) \leq D(E_1). \quad (12)$$

To see that f exists, note that by Proposition 2.2 there is a neighborhood of unity U in the group of linear transformation of \mathbb{P}^{i_N} such that for any $g \in U$ the morphism $g \circ \varphi_{1,N} : \mathbb{P}^1 \rightarrow \mathbb{P}^{i_N}$ (where $\varphi_{1,N} = \varphi_N \circ \varphi_{N-1} \circ \dots \circ \varphi_1$) satisfies (12). Moreover, it is obvious that one can choose g in such a way that (11) holds.

We now extend the morphism f to a morphism $\tilde{f} : \mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}^{i_N}$ in the following way. Let f be given in coordinates as $z_i = f_i(x_0, x_1)$. Identify x_0, x_1 with homogeneous coordinates on the second factor of $\mathbb{P}^1 \times \mathbb{P}^1$, and let t_0, t_1 be homogeneous coordinates on the first factor of $\mathbb{P}^1 \times \mathbb{P}^1$. Then define \tilde{f} by putting $z_i = t_0 f_i(x_0, x_1)$ for $i = 0, 1$ and $z_i = t_1 f_i(x_0, x_1)$ for $i \geq 2$. Condition (11) ensures that \tilde{f} is well-defined. Set $\tilde{E}_N := \tilde{f}_N^* E_N$.

The morphism \tilde{f} has the following properties.

a) The restriction of \tilde{f} to the vertical fiber P over the point $(t_0, t_1) = (1, 1)$ coincides with f . Therefore (12) implies

$$D(\tilde{E}_N|_P) \leq D(E_1). \quad (13)$$

b) The restriction of \tilde{f} to the vertical fiber P' over the point $(t_0, t_1) = (1, 0)$ is a morphism of P' onto l of degree $\deg \varphi_{1,N}$. Thus

$$D(\tilde{E}_N|_{P'}) = (\deg \varphi_{1,N}) D_N. \quad (14)$$

c) The restriction of \tilde{f} to any horizontal fiber S is a linear embedding. Therefore

$$D(\tilde{E}_N|_S) \leq D_N. \quad (15)$$

The hypotheses of Lemma 2.5 hold for \tilde{E}_N , since $c_1(\tilde{E}_N|_P) = c_1(\tilde{E}_N|_S) = 0$ by Lemma 3.2. Therefore Lemma 2.5, together with (13), (14), and (15), yields

$$(\deg \varphi_{1,N})D_N \leq 4(\text{rk } E_N)(D(E_1) + 2)(D_N + 1) - \chi(E_N) - \chi(E_N^*). \quad (16)$$

By the Riemann-Roch Theorem and the vanishing of all Chern classes of E_N (Lemma 3.2), we have

$$\chi(E_N) = \chi(E_N^*) = n\chi(\mathcal{O}_{P_N}) = \text{rk } E_N.$$

Thus (16) turns into

$$(\deg \varphi_{1,N})D_N \leq 4(\text{rk } E_N)(D(E_1) + 2)(D_N + 1) - 2 \text{rk } E_N. \quad (17)$$

As $\text{rk } E_N = \text{rk } E$ and $D(E_1)$ do not depend on N , we see that the right hand side of (17) is a linear function of D_N , while the left hand side grows with N faster than a linear function of D_N , since $\lim_{N \rightarrow \infty} \deg \varphi_{1,N} = \infty$ by our hypothesis. Therefore, inequality (17) can hold only if D_N equals zero for large enough N and hence for all N . This means that all vector bundles E_N are linearly trivial, and thus trivial. \square

Theorem 3.1 together with Theorem 2.1 leads to the following general description of vector bundles on an inductive limit of relative projective spaces. Let $p_N : M_N \rightarrow \mathcal{S}_N$ be a relative projective space, i.e. a locally trivial fibration with base a connected complex analytic space \mathcal{S}_N and with fiber the projective space \mathbb{P}^{i_N} . Let furthermore

$$\begin{array}{ccccc} M_1 & \xrightarrow{\varphi_1} & M_2 & \xrightarrow{\varphi_2} & \dots \\ p_1 \downarrow & & p_2 \downarrow & & \\ \mathcal{S}_1 & \longrightarrow & \mathcal{S}_2 & \longrightarrow & \dots \end{array} \quad (18)$$

be a commutative diagram. Note that if the restriction of φ_N to a fiber P_N of p_N has degree d , then the restriction of φ_N to any fiber of p_N has also degree d . Therefore it makes sense to call (18) *linear* if the degree of the restriction of φ_N on the fibers of p_N equals 1 for almost all N , and *twisted* otherwise.

Theorem 3.3. *Let $E = \{E_N\}$ be a vector bundle on the upper row of (18). Then there exist integers $d_1 \geq \dots \geq d_{\text{rk } E}$ such that, for large enough N and any fiber P_N of p_N , we have $E_N|_{P_N} \simeq \bigoplus_r \mathcal{O}_{P_N}(d_r)$. Furthermore, $d_1 = \dots = d_{\text{rk } E} = 0$ if (18) is twisted.*

Proof. The fact that $E_N|_{P_N}$ is isomorphic to a direct sum of line bundles follows from Theorems 2.1 and 3.1. The fact that the splitting data of $E_N|_{P_N}$ does not depend on P_N follows from Proposition 2.3. If (18) is linear, the degree of P_N equals 1 for large enough N , thus the splitting data $E_N|_{P_N}$ also does not depend on N for large enough N . Finally, if (18) is twisted, each $d_r(E_N|_{P_N})$ is divisible by $(\deg \varphi_N) \cdot \dots \cdot (\deg \varphi_j)$ for all $j > N$, and therefore $d_r(E_N|_{P_N}) = 0$. \square

It is not true in general that E is isomorphic to a direct sum of line bundles, or that E admits a filtration whose associated quotients are line bundles. To see this it suffices to consider the example when $\mathcal{S} = \mathcal{S}_N = \mathbb{P}^2$ for each N and E is the pullback of a rank 2 bundle $E_{\mathcal{S}}$ on \mathbb{P}^2 with no line subbundle, for instance the tautological bundle S_2 . However, the reader will easily prove that, if all d_r are distinct (that is $\nu_r = 1$ for all r), E always admits a filtration whose associated quotients are line bundles. Finally, if (18) is twisted, Theorem 3.3 implies that each E_N is the pull-back $p_N^* E'_N$ for some vector bundle $E' = \{E'_N\}$ on the lower row of (18).

4 The case of ind-grassmannians

In this section we consider two different types of closed immersions

$$G(k_1, n_1) \xrightarrow{f_1} G(k_2, n_2) \xrightarrow{f_2} \dots \quad (19)$$

and characterize vector bundles on the corresponding ind-spaces.

4.1 Standard extensions of grassmannians

We define a *standard extension* of grassmannians as a closed immersion of the form

$$\lambda_{r,m} : G(k, n) \rightarrow G(k+r, n+m), \quad \{V \subset \mathbb{C}^n\} \mapsto \{V \oplus W \subset \mathbb{C}^n \oplus \mathbb{C}^m\},$$

where $W \subset \mathbb{C}^m$ is a fixed subspace of dimension $r \geq 0$.

Proposition 4.1. *Assume that a sequence (19) is given, where f_N are standard extensions. Let $E = \{E_N\}$ be a vector bundle on (19). Then for each N , E_N is a uniform bundle on $G(k_N, n_N)$.*

Proof. For each N consider the natural diagram

$$\begin{array}{ccc} F(k_N - 1, k_N, n_N) & \xrightarrow{\pi_N} & G(k_N, n_N) \\ p_N \downarrow & & \\ G(k_N - 1, n_N) & & \end{array} \quad (20)$$

where $F(k_N - 1, k_N, n_N)$ stands for the space of all flags of type $(k_N - 1, k_N)$ in \mathbb{C}^{n_N} . Note that f_N induce morphisms between the diagrams (20) for N and $N + 1$. Furthermore, the embeddings $F(k_N - 1, k_N, n_N) \rightarrow F(k_{N+1} - 1, k_{N+1}, n_{N+1})$ and the morphisms p_N define a relative projective ind-space. The vector bundles $\pi_N^* E_N$ define a vector bundle on this relative projective ind-space. By Theorem 3.3 the splitting data of the restriction $\pi_N^* E_N|_{P_N}$ on each fiber P_N of p_N does not depend on N and P_N . This implies the result, as any projective line $l \subset G(k_N, n_N)$ is a line in some fiber P_N , and the splitting data of $E|_l$ and $\pi_N^* E_N|_{P_N}$ are equal. \square

Theorem 4.2. *Assume that a sequence (19) is given, where f_N are standard extensions and $\lim_{N \rightarrow \infty} k_N = \lim_{N \rightarrow \infty} (n_N - k_N) = \infty$. Then any vector bundle $E = \{E_N\}$ on the inductive limit of (19) is isomorphic to a direct sum of line bundles.*

Proof. By Proposition 4.1, each E_N is a uniform bundle. We will prove the Theorem by induction on $\text{rk } E = \text{rk } E_N$. Fix N and let $d_1, \dots, d_{\text{rk } E_N}$ be the splitting data for the restriction $E_N|_l$, where $l \subset G(k_N, n_N)$ is any line. Denote by ν_1 the multiplicity of d_1 . As each f_j induces an isomorphism of Picard groups, the splitting data $d_1, \dots, d_{\text{rk } E_N}$ does not depend on N . Fix $x \in G(k_N, n_N)$. Any line $l \subset G(k_N, n_N)$ passing through x determines a subspace $E'_{l,x}$ in the fiber $(E_N)_x$: this is the fiber of the unique subbundle of $E_N|_l$ isomorphic to the direct sum of ν_1 copies of $\mathcal{O}_l(d_1)$, see Subsection 2.2. The variety of all lines l passing through x is isomorphic to the direct product $\mathbb{P}^{k_N-1} \times \mathbb{P}^{n_N-k_N-1}$, hence we have a morphism

$$\psi_x : \mathbb{P}^{k_N-1} \times \mathbb{P}^{n_N-k_N-1} \longrightarrow G(\nu_1, \text{rk } E_N), \quad l \mapsto E'_{l,x}.$$

By our assumption, $\lim_{N \rightarrow \infty} k_N = \lim_{N \rightarrow \infty} (n_N - k_N) = \infty$. Thus, for large N , the morphism ψ_x is trivial, i.e. it determines a fixed subspace $(E'_N)_x$ in $(E_N)_x$. In this way we obtain a subbundle $E'_N \subset E_N$. Clearly $f_N^*(E'_{N+1}) = E'_N$, therefore $E' := \{E'_N\}$ is a well-defined subbundle of E .

If $\text{rk } E' = \text{rk } E$, twisting by the line bundle $\{\mathcal{O}_{G(k_N, n_N)}(-d_1)\}$ yields a bundle whose restriction to any line in $G(k_N, n_N)$ is trivial for all N . Therefore the twisted bundle is trivial on each $G(k_N, n_N)$, and hence trivial. Thus E is isomorphic to a direct sum of ν_1 copies of $\{\mathcal{O}_{G(k_N, n_N)}(d_1)\}$. If $\text{rk } E' < \text{rk } E$, the induction assumption implies that both E' and E/E' are isomorphic to direct sums of line bundles. Finally, the Bott-Borel-Weil Theorem implies that $H^1(\mathcal{L}) = 0$ for any line bundle \mathcal{L} on $G(k_N, n_N)$ unless $k_N = n_N - k_N = 1$. (More precisely, the equality $H^1(\mathcal{L}) = 0$ follows from Bott's Vanishing Theorem, see [B] and [De], via the observation that $H^1(\mathcal{L}) = H^1(\mathcal{L}')$, where \mathcal{L}' is the pullback of \mathcal{L} on the full flag variety). Therefore, there are no non-trivial extensions of line bundles on $G(k_N, n_N)$ for $k_N > 1$. This implies that E is isomorphic to a direct sum of line bundles. \square

Here is an important special case of Theorem 4.2. If $V \subset \mathbb{C}^\infty$ is an arbitrary fixed subspace, a subspace $V' \subset \mathbb{C}^\infty$ is *commensurable* with V , if there exists a finite dimensional subspace $U \subset \mathbb{C}^\infty$ such that $V \subset V' + U$, $V' \subset V + U$, and $\dim V \cap U = \dim V' \cap U$. The set of all subspaces V' comensurable with V is, by definition, the ind-grassmannian $G(V, \infty)$. If V is finite dimensional and $\dim V = k$, then $G(V, \infty) = G(k, \infty)$. In [DP] an explicit construction of $G(V, \infty)$ as an ind-space is given. Moreover, $G(V, \infty)$ is the inductive limit of standard extensions, and the conditions of Theorem 4.2 are satisfied for $G(V, \infty)$ if and only if $\dim V = \text{codim}_{\mathbb{C}^\infty} V = \infty$. Therefore, by Theorem 4.2, we conclude that in the latter case every vector bundle on $G(V, \infty)$ is isomorphic to a direct sum of line bundles.

More generally, if $V_1 \subset \dots \subset V_r$ is any fixed flag in \mathbb{C}^∞ , the ind-variety $F(V_1, \dots, V_r, \infty)$ of flags $V'_1 \subset \dots \subset V'_r$ comensurable with $V_1 \subset \dots \subset V_r$, is constructed in [DP]. We leave it to the reader to prove the following corollary of Theorem 4.2 by double induction on $\text{rk } E$ and r .

Corollary 4.3. *Let $\dim V_1 = \dim(V_2/V_1) = \dots = \dim(\mathbb{C}^\infty/V_r) = \infty$. Then every vector bundle E on $F(V_1, \dots, V_r, \infty)$ is isomorphic to a direct sum of line bundles.*

4.2 Twisted extensions of grassmannians

An alternative definition of a grassmannian is as a homogeneous space $GL(n)/P$ for a maximal parabolic subgroup $P \subset GL(n)$. We call a morphism $f : G(k_1, n_1) \rightarrow G(k_2, n_2)$ *homogeneous* if it is induced by a group homomorphism $\tilde{f} : GL(n_1) \rightarrow GL(n_2)$. A homogeneous morphism is either a closed immersion or its image is a point.

Consider a homogeneous morphism $f : G(k_1, n_1) \rightarrow G(k_2, n_2)$, where $G(k_1, n_1) = GL(n_1)/P_1$, $G(k_2, n_2) = GL(n_2)/P_2$, and P_1, P_2 are maximal parabolic subgroups such that $\tilde{f}(P_1) \subset P_2$. The reductive part of P_1 is isomorphic to $GL(k_1) \times GL(n_1 - k_1)$ and the reductive part of P_2 is isomorphic to $GL(k_2) \times GL(n_2 - k_2)$. Note that $\tilde{f}(GL(k_1) \times GL(n_1 - k_1))$ is contained in one component of the direct product $GL(k_2) \times GL(n_2 - k_2)$ if and only if the induced morphism $f : G(k_1, n_1) \rightarrow G(k_2, n_2)$ is a morphism into a point. In the sequel we adopt a convention: whenever we write a homogeneous morphism as $f : G(k_1, n_1) \rightarrow G(k_2, n_2)$ and f is a closed immersion, we automatically assume that $\tilde{f}(GL(k_1)) \subset GL(k_2)$ and $\tilde{f}(GL(n_1 - k_1)) \subset GL(n_2 - k_2)$. Furthermore, we call a morphism $\varphi : \mathbb{P}^N \rightarrow G(k, n)$ *k-split* if φ^*S_k is isomorphic to a direct sum of line bundles on \mathbb{P}^N . For example, Theorem 2.1 implies that any morphism $\mathbb{P}^1 \rightarrow G(k, n)$ is *k-split*.

Proposition 4.4. *a) Any homogeneous morphism $\varphi : \mathbb{P}^N = G(1, N) \rightarrow G(k, n)$ is *k-split*.*

b) Let $f : G(k_1, n_1) \rightarrow G(k_2, n_2)$ be a homogeneous embedding of grassmannians and $\varphi : \mathbb{P}^N \rightarrow G(k_1, n_1)$ be a k_1 -split morphism. Then the composition $f \circ \varphi : \mathbb{P}^N \rightarrow G(k_2, n_2)$ is k_2 -split.

Proof. a) Let $\mathbb{P}^N = GL(N+1)/P_1$, and $G(k, N) = GL(N)/P_2$. The reductive parts of P_1 and P_2 are isomorphic respectively to $GL(1) \times GL(N)$ and $GL(k) \times GL(N-k)$. According to our convention, the group homomorphism $\tilde{\varphi} : GL(N+1) \rightarrow GL(N)$ which induces φ maps $GL(1)$ into $GL(k)$, and $GL(N)$ into $GL(N-k)$. Therefore the structure group of φ^*S_k is reduced to $GL(1)$, which implies that φ^*S_k is a direct sum of line bundles.

b) Let $\tilde{f} : GL(n_1) \rightarrow GL(n_2)$ be a group homomorphism which induces the morphism f . Since $\tilde{f}(GL(k_1)) \subset GL(k_2)$, the $GL(n_1)$ -homogeneous bundle $f^*S_{k_2}$ is a direct summand in the tensor product of several copies of S_{k_1} and $S_{k_1}^*$. This, together with the fact, that the morphism φ is k_1 -split, implies that $(f \circ \varphi)^*S_{k_2}$ is a direct summand of a bundle isomorphic to a direct sum of line bundles on \mathbb{P}^N . Hence $(f \circ \varphi)^*S_{k_2}$ is isomorphic itself to a direct sum of line bundles, i.e. the morphism $f \circ \varphi$ is k_2 -split. \square

Here is a coordinate form of a *k-split* map. Let $t = (t_0, \dots, t_N)$ be homogeneous coordinates on \mathbb{P}^N , and v_1, \dots, v_n be a basis in \mathbb{C}^n . Then the reader will check that a morphism $\varphi : \mathbb{P}^N \rightarrow G(k, n)$ is *k-split* if and only if φ can be presented in the form

$$\varphi(t) = \text{span} \left\{ \sum_{j=1}^n \varphi_{i,j}(t) v_j, \quad i = 1, \dots, k \right\} \quad (21)$$

for some homogeneous polynomials $\varphi_{i,j}(t)$ in t . It is clear that if $\varphi : \mathbb{P}^N \rightarrow G(k, n)$ is represented in the form (21), then $\deg \varphi_{i,j}$ does not depend on j and $\deg \varphi = \sum_{i=1}^k \deg \varphi_{i,j}$.

Lemma 4.5. *Let $\varphi : \mathbb{P}^N \rightarrow G(k, n)$ be a k -split morphism and $\lambda_{0,k} : G(k, n) \rightarrow G(k, n+k)$ be a standard extension (see Subsection 4.1). Then the composition $\lambda_{0,k} \circ \varphi : \mathbb{P}^N \rightarrow G(k, n+k)$ factors through a standard extension $\mathbb{P}^N \rightarrow \mathbb{P}^{N+1}$, via a k -split morphism $\tilde{\varphi} : \mathbb{P}^{N+1} \rightarrow G(k, n+k)$.*

Proof. Let φ be presented in the form (21), and let the vectors v_{n+1}, \dots, v_{n+k} extend the set v_1, \dots, v_n to a basis in \mathbb{C}^{n+k} . Fix homogeneous coordinates $t' = (t_0, \dots, t_{N+1})$ in \mathbb{P}^{N+1} and identify \mathbb{P}^N with the hyperplane $t_{N+1} = 0$. Then $\tilde{\varphi} : \mathbb{P}^{N+1} \rightarrow G(k, n+k)$ is given by the formula

$$\tilde{\varphi}(t') = \text{span} \left\{ \sum_{j=1}^n \varphi_{i,j}(t) v_j + t_{N+1}^{\delta_i} v_{n+i}, \quad i = 1, \dots, k \right\},$$

where $\delta_i := \deg \varphi_{i,j}$. □

We need one last definition. We call a homogeneous morphism $f : G(k_1, n_1) \rightarrow G(k_2, n_2)$ a *twisted extension* if f decomposes as $\lambda_{0, n_2 - n_1} \circ h$, where $h : G(k_1, n_1) \rightarrow G(k_2, n_1)$ is a homogeneous morphism for $n_2 - n_1 \geq k_2$, and $\lambda_{0, n_2 - n_1} : G(k_2, n_1) \rightarrow G(k_2, n_2)$ is a standard extension.

Proposition 4.6. *Assume that a sequence (19) is given, where f_N are twisted extensions, infinitely many of which have degree greater than 1. Suppose in addition that $\lim_{N \rightarrow \infty} k_N = \lim_{N \rightarrow \infty} (n_N - k_N) = \infty$. Then any vector bundle $E = \{E_N\}$ on the inductive limit of (19) is trivial.*

Proof. Let $E = \{E_N\}$ be a vector bundle on the inductive limit of (19). Fix N and a $(k_N - 1)$ -dimensional subspace of \mathbb{C}^{n_N} . This determines a projective subspace $\mathbb{P}^{n_N - k_N}$ of $G(k_N, n_N)$, and the closed immersion $\varphi_N : \mathbb{P}^{n_N - k_N} \rightarrow G(k_N, n_N)$ is k_N -split as $\varphi_N^* S_{k_N} \simeq (k_N - 1) \mathcal{O}_{\mathbb{P}^{n_N - k_N}} \oplus \mathcal{O}_{\mathbb{P}^{n_N - k_N}}(-1)$. Furthermore, let $f_N = \lambda_{0, m_N} \circ h_N$, where $h_N : G(k_N, n_N) \rightarrow G(k_{N+1}, n_{N+1} - m_N)$ is a homogeneous closed immersion and $m_N \geq k_{N+1}$. Proposition 4.4 b) implies that the composition $h_N \circ \varphi_N$ is k_{N+1} -split. Then, by Lemma 4.5, the morphism $f_N \circ \varphi_N = \lambda_{0, m_N} \circ (h_N \circ \varphi_N)$ factors through a standard extension $\mathbb{P}^{n_N - k_N} \rightarrow \mathbb{P}^{n_N - k_N + 1}$ via a homogeneous immersion $\varphi_{N+1} : \mathbb{P}^{n_N - k_N + 1} \rightarrow G(k_{N+1}, n_{N+1})$. Proceeding by induction on j , we build homogeneous immersions $\varphi_{N+j} : \mathbb{P}^{n_N - k_N + j} \rightarrow G(k_{N+j}, n_{N+j})$ and standard extensions $\mathbb{P}^{n_N - k_N + j} \rightarrow \mathbb{P}^{n_N - k_N + j + 1}$ which form the commutative diagram

$$\begin{array}{ccccccc} \mathbb{P}^{n_N - k_N} & \longrightarrow & \mathbb{P}^{n_N - k_N + 1} & \longrightarrow & \dots & & \\ \varphi_N \downarrow & & \varphi_{N+1} \downarrow & & & & \\ G(k_N, n_N) & \xrightarrow{f_N} & G(k_{N+1}, n_{N+1}) & \xrightarrow{f_{N+1}} & \dots & & \end{array} \quad (22)$$

Applying the BVT theorem to the inductive limit of the upper row of (22), we conclude that $E_N|_{\mathbb{P}^{n_N - k_N}}$ is isomorphic to a direct sum of line bundles. Furthermore, one sees exactly as in Proposition 4.1 that E_N is a uniform bundle. Now the argument in the

proof of Theorem 4.2 shows that, since $\lim_{N \rightarrow \infty} k_N = \lim_{N \rightarrow \infty} (n_N - k_N) = \infty$, E_N is isomorphic to a direct sum of line bundles. Therefore E is also isomorphic to a direct sum of line bundles. Finally, as $\deg f_N > 1$ for infinitely many N , a line bundle on this inductive limit is necessarily trivial, hence E is trivial. \square

Here are two examples to Proposition 4.6.

Let \mathbb{C}^w be a proper subspace of \mathbb{C}^m . Consider the closed immersion

$$G(k, n) \longrightarrow G(kw, nm), \quad \{V \subset \mathbb{C}^n\} \mapsto \{V \otimes \mathbb{C}^w \subset \mathbb{C}^n \otimes \mathbb{C}^m\}. \quad (23)$$

It factors, via a standard extension, through the closed immersion

$$G(k, n) \longrightarrow G(kw, nw), \quad \{V \subset \mathbb{C}^n\} \mapsto \{V \otimes \mathbb{C}^w \subset \mathbb{C}^n \otimes \mathbb{C}^w\}.$$

Fixing a sequence of pairs $\mathbb{C}^{w_N} \subset \mathbb{C}^{m_N}$, determines a sequence of immersions (23). If the set $\{w_N\}$ is bounded, for sufficiently large N all embeddings in this sequence are twisted extensions as

$$nm_1 \dots m_{N+1} - nw_1 \dots w_{N+1} \geq kw_1 \dots w_{N+1}$$

for large enough N . Therefore Proposition 4.6 implies that any vector bundle on the inductive limit is trivial.

Finally consider a closed immersion of the form

$$G(k, n) \longrightarrow G(k(k+1)/2, n^2), \quad \{V \subset \mathbb{C}^n\} \mapsto \{S^2(V) \subset \mathbb{C}^n \otimes \mathbb{C}^n\}, \quad (24)$$

where S^2 stands for symmetric square. By iteration of (24) we obtain another example of a sequence of twisted extensions. Indeed, (24) factors, via a standard extension, through the closed immersion

$$G(k, n) \longrightarrow G(k(k+1)/2, n(n+1/2)), \quad \{V \subset \mathbb{C}^n\} \mapsto \{S^2(V) \subset S^2(\mathbb{C}^n)\},$$

and the inequality

$$k(k+1)/2 \leq n^2 - n(n+1)/2$$

needed in the definition of a twisted extension, is trivially satisfied. Proposition 4.6 implies that any vector bundle on the corresponding inductive limit is also trivial.

5 Conclusion

The results of this paper lead naturally to the following question. Consider an arbitrary sequence of morphisms

$$G(k_1, n_1) \xrightarrow{f_1} G(k_2, n_2) \xrightarrow{f_2} \dots \quad (25)$$

Is it true that, if $E = \{E_N\}$ is an arbitrary vector bundle on (25) then each E_N is a homogeneous bundle on $G(k_N, n_N)$? In fact, the BVT Theorem, both main theorems of

Sato (see [S2]), Theorems 3.1 and 4.2, as well as Proposition 4.6, are all equivalent to the affirmative answer to the above question for the cases they apply to. More precisely, the description of vector bundles on sequences (25) in each specific case considered in the above statements, is equivalent to the descriptions of systems of homogeneous bundles $\{E_N\}$ with $f_{N+1}^*E_{N+1} = E_N$. Note that Theorem 3.1 applies to not necessarily homogeneous morphisms, while in all other statements homogeneous morphisms are considered. It would be very interesting to give an answer to the above general question even under the assumption that all morphisms f_N are homogeneous.

References

- [B] R. Bott, Homogeneous vector bundles, *Ann. of Math* (2) **66** (1957), 203-248.
- [BV] W. Barth and A. Van de Ven, On the geometry in codimension 2 in Grassmann manifolds, *Lecture Notes in Math.* 412 (1974), 1–35.
- [De] M. Demazure, Une démonstration algébrique d’un théorème de Bott (French), *Invent. Math.* **5** (1968), 349-356.
- [DPW] I. Dimitrov, I. Penkov, and J. A. Wolf, A Bott-Borel-Weil theorem for direct limits of algebraic groups, *Amer. J. of Math.* **124** (2002), 955–998.
- [DP] I. Dimitrov and I. Penkov, Ind-varieties of generalized flags as homogeneous spaces for classical ind-groups, preprint 2001.
- [Do1] J. Donin, Deformations of holomorphic vector bundles on the Riemann sphere, *Mat. Sb. (N.S.)* **71(113)** (1966), 495–502 (Russian).
- [Do2] J. Donin, Construction of a versal family of deformations for holomorphic principal bundles over a compact complex space, *Mat. Sb. (N.S.)* **94(136)** (1974), 430–443 (Russian).
- [P] I. Penkov, The Penrose transform on general grassmannians, *C. R. Acad. Bulgare Sci.* **33** (1980), 1439–1442.
- [S1] E. Sato, On the decomposability of infinitely extendable vector bundles on projective spaces and Grassmann varieties, *J. Math. Kyoto Univ.* **17** (1977), 127–150.
- [S2] E. Sato, On infinitely extendable vector bundles on G/P , *J. Math. Kyoto Univ.* **19** (1979), 171–189.
- [T] A. N. Tyurin, Vector bundles of finite rank over infinite varieties, *Math. USSR Izvestija*, **10** (1976), 1187–1204.

J. Donin
 Department of Mathematics
 Bar-Ilan University
 52900 Ramat Gan, Israel
 donin@macs.biu.ac.il

I. Penkov
 Department of Mathematics

University of California Riverside
Riverside, CA 92521, USA
penkov@math.ucr.edu