

BOUNDED SIMPLE $(\mathfrak{g}, \mathfrak{sl}(2))$ -MODULES FOR $\operatorname{rk} \mathfrak{g} = 2$

IVAN PENKOV AND VERA SERGANOVA

ABSTRACT. This paper is a continuation of our work [PS2] in which we prove some general results about simple $(\mathfrak{g}, \mathfrak{k})$ -modules with bounded \mathfrak{k} -multiplicities (or bounded simple $(\mathfrak{g}, \mathfrak{k})$ -modules). In the absence of a classification of bounded simple $(\mathfrak{g}, \mathfrak{k})$ -modules in general, it is important to understand some special cases as best as possible. Here we consider the case $\mathfrak{k} = \mathfrak{sl}(2)$. It turns out that in order for an infinite-dimensional bounded simple $(\mathfrak{g}, \mathfrak{sl}(2))$ -module to exist, \mathfrak{g} must have rank 2, and, up to conjugation, there are five possible embeddings $\mathfrak{sl}(2) \rightarrow \mathfrak{g}$ which yield infinite-dimensional bounded simple $(\mathfrak{g}, \mathfrak{sl}(2))$ -modules.

Our main result is a detailed description of the bounded simple $(\mathfrak{g}, \mathfrak{sl}(2))$ -modules in all five cases. When $\mathfrak{g} \simeq \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$ we reproduce in modern terms some classical results from the 1940's. When $\mathfrak{g} \simeq \mathfrak{sl}(3)$ and $\mathfrak{sl}(2)$ is a principal subalgebra, bounded simple $(\mathfrak{sl}(3), \mathfrak{sl}(2))$ -modules are Harish-Chandra modules and our result singles out all Harish-Chandra modules with bounded $\mathfrak{sl}(2)$ -multiplicities. A case where the result is entirely new is the case of a principal $\mathfrak{sl}(2)$ -subalgebra of $\mathfrak{g} = \mathfrak{sp}(4)$.

1. INTRODUCTION

The classification of simple Harish-Chandra modules is a celebrated result and there is an extensive literature on the general topic of Harish-Chandra modules, see for instance [KV] and the references therein. Algebraically, Harish-Chandra modules are $(\mathfrak{g}, \mathfrak{k})$ -modules for a symmetric subalgebra \mathfrak{k} of a semisimple Lie algebra \mathfrak{g} , and in the last decade an intense exploration of more general $(\mathfrak{g}, \mathfrak{k})$ -modules for not necessarily symmetric subalgebras \mathfrak{k} has begun, [PS1],[PSZ], [PZ1], [PZ2], [PZ3]. A most notable result in this direction is the classification of simple $(\mathfrak{g}, \mathfrak{k})$ -modules of finite type and a generic minimal \mathfrak{k} -type carried out in [PZ2]. Nevertheless, the classification problem for $(\mathfrak{g}, \mathfrak{k})$ -modules of finite type with an arbitrary minimal \mathfrak{k} -type is still open even when $\operatorname{rk} \mathfrak{g} = 2$ and $\operatorname{rk} \mathfrak{k} = 1$.

2000 *Mathematics Subject Classification*. Primary 17B10, Secondary 22E46.

In the recent paper [PS2] we concentrated on the interesting subclass of bounded $(\mathfrak{g}, \mathfrak{k})$ -modules (the definition see in Section 1 below) and proved some general results regarding the existence of such modules. In particular, we established sufficient and necessary conditions on a reductive in \mathfrak{g} subalgebra \mathfrak{k} for the existence of a simple infinite-dimensional bounded $(\mathfrak{g}, \mathfrak{k})$ -module. If $\mathfrak{k} \simeq \mathfrak{sl}(2)$, simple $(\mathfrak{g}, \mathfrak{sl}(2))$ -modules of finite type exist for any simple \mathfrak{g} , see for instance [PSZ] or [PZ3]. It turns out however, that bounded $(\mathfrak{g}, \mathfrak{sl}(2))$ -modules are very special and exist only for $\mathrm{rk} \mathfrak{g} = 2$. The classification of such modules is rather intriguing as they are the "smallest", and thus highly non-generic, $(\mathfrak{g}, \mathfrak{sl}(2))$ -modules.

This classification is carried out in the present paper. We show first that, up to conjugation, there are precisely five possibilities for embedding $\mathfrak{sl}(2)$ into a Lie algebra of \mathfrak{g} of rank 2 so that bounded infinite-dimensional $(\mathfrak{g}, \mathfrak{sl}(2))$ -modules exist: $\mathfrak{sl}(2)$ as the diagonal subalgebra of $\mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$, $\mathfrak{sl}(2)$ as a root subalgebra or a principal $\mathfrak{sl}(2)$ subalgebra of $\mathfrak{sl}(3)$, and $\mathfrak{sl}(2)$ as a root subalgebra corresponding to a short root or as a principal subalgebra of $\mathfrak{sp}(4)$.

We then give a classification and a detailed description (we compute characters and minimal $\mathfrak{sl}(2)$ -types) of all bounded $(\mathfrak{g}, \mathfrak{sl}(2))$ -modules. In the case when $\mathfrak{g} \simeq \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$ our results are just a modern reproduction of classical results, in all other cases they are new. The most interesting new case is that of a principal $\mathfrak{sl}(2)$ -subalgebra of $\mathfrak{sp}(4)$.

Acknowledgement. This paper has been written in close contact with Gregg Zuckerman who has supported us on several occasions with valuable advice. David Vogan, Jr. has also generously shared his knowledge of Harish-Chandra modules with us. We thank T. Milev for reading the manuscript carefully and checking some of the calculations. Finally, we acknowledge the hospitality and support of the Max Planck Institute for Mathematics in Bonn.

2. GENERAL DEFINITIONS AND PRELIMINARY RESULTS

The ground field is \mathbb{C} .

Let \mathfrak{g} be a semisimple (finite-dimensional) Lie algebra and $\mathfrak{k} \subset \mathfrak{g}$ be a reductive in \mathfrak{g} subalgebra. $U(\cdot)$ stands for enveloping algebra, $U = U(\mathfrak{g})$ and Z_U is the center of U . A $(\mathfrak{g}, \mathfrak{k})$ -module M is a \mathfrak{g} -module M on which \mathfrak{k} acts locally finitely, i.e. $\dim U(\mathfrak{k}) \cdot m < \infty$, $\forall m \in M$. A $(\mathfrak{g}, \mathfrak{k})$ -module M has *finite type over \mathfrak{k}* if the Jordan–Hölder multiplicity of any fixed simple finite-dimensional \mathfrak{k} -module V (such a V is called a \mathfrak{k} -type) in arbitrary finite-dimensional \mathfrak{k} -submodules of M is bounded. A $(\mathfrak{g}, \mathfrak{k})$ -module is *bounded* if the above multiplicities are bounded by a

constant not depending on the \mathfrak{k} -type V . A reductive in \mathfrak{g} subalgebra $\mathfrak{k} \subset \mathfrak{g}$ is *bounded* if there exists an infinite-dimensional simple bounded $(\mathfrak{g}, \mathfrak{k})$ -module M . A bounded subalgebra $\mathfrak{k} \subset \mathfrak{g}$ is *strictly bounded* if there is a simple bounded $(\mathfrak{g}, \mathfrak{k})$ -module M on which no simple ideal of \mathfrak{g} acts locally finitely. The following necessary conditions on a subalgebra \mathfrak{k} to be bounded, or strictly bounded, are proved in [PS2] (Theorem 4.1 and Corollary 4.6).

Theorem 2.1. *Let \mathfrak{k} be a bounded reductive subalgebra of a semisimple Lie algebra $\mathfrak{g} = \bigoplus_i \mathfrak{g}_i$ (\mathfrak{g}_i being the simple ideals of \mathfrak{g}).*

a) *If M is a simple bounded $(\mathfrak{g}, \mathfrak{k})$ -module and the algebra of \mathfrak{k} -invariants $\mathfrak{g}_{i_0}^{\mathfrak{k}}$ is not abelian for some i_0 , then $M \simeq M_{i_0} \otimes \overline{M_{i_0}}$, where M_{i_0} is a simple finite-dimensional $\mathfrak{g}_{i_0}^{\mathfrak{k}}$ -module and $\overline{M_{i_0}}$ is a simple bounded $(\bigoplus_{i \neq i_0} \mathfrak{g}_i, (\bigoplus_{i \neq i_0} \mathfrak{g}_i) \cap \mathfrak{k})$ -module.*

b) *If $r_{\mathfrak{g}}$ is the half-dimension of a nilpotent orbit of minimal positive dimension in \mathfrak{g} , then*

$$(1) \quad r_{\mathfrak{g}} \leq b_{\mathfrak{k}},$$

where $b_{\mathfrak{k}}$ is the dimension of a Borel subalgebra of \mathfrak{k} .

c) *If \mathfrak{k} is strictly bounded, then*

$$\sum_i r_{\mathfrak{g}_i} \leq b_{\mathfrak{k}}.$$

In [PS2] we also established the following sufficient condition for a reductive in \mathfrak{g} subalgebra $\mathfrak{k} \subset \mathfrak{g}$ to be bounded. Recall that a finite-dimensional module W over an algebraic group H is *spherical* if a Borel subgroup B_H has an open orbit in W .

Theorem 2.2. *Let $K \subset G \subset GL(V)$ be a chain of reductive algebraic groups, and let $V' \subset V$ be a 1-dimensional space whose stabilizers in G and K are parabolic subgroups $P \subset G$ and $Q \subset K$. Then, if $(V')^* \otimes (\mathfrak{g} \cdot V' / \mathfrak{k} \cdot V')$ is a spherical module over a reductive part Q_0 of Q , \mathfrak{k} is a bounded subalgebra of \mathfrak{g} .*

3. BOUNDED SUBALGEBRAS OF A RANK-2 LIE ALGEBRA

Our main interest in this paper are infinite-dimensional bounded $(\mathfrak{g}, \mathfrak{k})$ -modules for $\mathfrak{k} \simeq \mathfrak{sl}(2)$. Theorem 1 implies that if $\mathfrak{k} \simeq \mathfrak{sl}(2)$ is a strictly bounded subalgebra of $\mathfrak{g} = \bigoplus_i \mathfrak{g}_i$, then $\sum_i r_{\mathfrak{g}_i} \leq 2$. This is easily seen to imply $\text{rk}\mathfrak{g} = 2$. Therefore, in the rest of the paper we restrict ourselves to the case when $\text{rk}\mathfrak{g} = 2$. The following theorem classifies more generally all reductive in \mathfrak{g} bounded subalgebras $\mathfrak{k} \subset \mathfrak{g}$ under the assumption that $\text{rk}\mathfrak{g} = 2$.

Theorem 3.1. *Let \mathfrak{g} be a semisimple Lie algebra of rank 2 and $\mathfrak{k} \subset \mathfrak{g}$ be a reductive in \mathfrak{g} bounded subalgebra. The following is a complete list of such pairs up to conjugation by inner automorphisms.*

- (1) $\mathfrak{g} \simeq \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$: $\mathfrak{k} \simeq \mathfrak{gl}(2)$ is a direct sum of a simple ideal and a Cartan subalgebra of the other simple ideal, $\mathfrak{k} \simeq \mathfrak{sl}(2)$ is a diagonal subalgebra, or \mathfrak{k} is any non-trivial toral subalgebra;
- (2) $\mathfrak{g} \simeq \mathfrak{sl}(3)$: \mathfrak{k} is a root subalgebra isomorphic to $\mathfrak{sl}(2)$ or $\mathfrak{gl}(2)$, \mathfrak{k} is a principal $\mathfrak{sl}(2)$ -subalgebra, or \mathfrak{k} is a Cartan subalgebra;
- (3) $\mathfrak{g} \simeq \mathfrak{sp}(4)$: $\mathfrak{k} \simeq \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$ is the subalgebra generated by the long roots, $\mathfrak{k} \simeq \mathfrak{gl}(2)$ is any root subalgebra, $\mathfrak{k} \simeq \mathfrak{sl}(2)$ is a root subalgebra corresponding to a short root, \mathfrak{k} is a principal $\mathfrak{sl}(2)$ -subalgebra, or \mathfrak{k} is a Cartan subalgebra;
- (4) $\mathfrak{g} \simeq G_2$: \mathfrak{k} is any subalgebra containing a Cartan subalgebra, in this case $\mathfrak{k} \simeq \mathfrak{sl}(3)$, $\mathfrak{k} \simeq \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$, or $\mathfrak{k} \simeq \mathfrak{gl}(2)$.

Proof. The inequality (1) implies that a 1-dimensional toral subalgebra is not bounded in all cases but (1). In (1) any 1-dimensional toral subalgebra \mathfrak{t} is bounded as the outer tensor product of a Verma module over a suitable ideal of \mathfrak{g} with the trivial module of the complementary ideal of \mathfrak{g} is always bounded as a $(\mathfrak{g}, \mathfrak{t})$ -module.

Similarly, (1) implies that a Cartan subalgebra is not bounded in G_2 . In all other cases it is well known to be bounded, see for instance [F].

If $\mathfrak{k} \simeq \mathfrak{sl}(2)$ then \mathfrak{k} is not bounded in G_2 again by (1), and if \mathfrak{k} is an ideal of $\mathfrak{g} = \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$, it is not bounded by Theorem 1 a). Furthermore, if $\mathfrak{k} \simeq \mathfrak{sl}(2)$ is a root subalgebra of $\mathfrak{g} = \mathfrak{sp}(4)$ corresponding to a long root, then \mathfrak{k} is not bounded by Theorem 1 a). For the remaining five possible embeddings of $\mathfrak{sl}(2)$ into a Lie algebra of rank 2, the image \mathfrak{k} is always a bounded subalgebra. This follows for instance from the explicit description of bounded $(\mathfrak{g}, \mathfrak{k})$ -modules which we present in Sections 4-7 of this paper.

For any embedding of $\mathfrak{gl}(2)$ into a Lie algebra \mathfrak{g} of rank 2, $\mathfrak{g} \not\cong G_2$, any generalized Verma module, corresponding to a parabolic subalgebra \mathfrak{p} which contains the image \mathfrak{k} of $\mathfrak{gl}(2)$, is a bounded $(\mathfrak{g}, \mathfrak{k})$ -module.

Consider next the case $\mathfrak{k} \simeq \mathfrak{sl}(2) \oplus \mathfrak{sl}(2) \subset \mathfrak{g}$ for $\mathfrak{g} = \mathfrak{sp}(4)$ or G_2 . Here the pair $(\mathfrak{g}, \mathfrak{k})$ is symmetric. In [V1] and [V2] ladder $(\mathfrak{g}, \mathfrak{k})$ -modules are constructed. Fix a Borel subalgebra $\mathfrak{b}_{\mathfrak{k}} \subset \mathfrak{k}$. By definition, a ladder module M has the \mathfrak{k} decomposition $M = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} V_{\mu+n\beta}$, where μ is some integral $\mathfrak{b}_{\mathfrak{k}}$ -dominant weight and β is the $\mathfrak{b}_{\mathfrak{k}}$ -highest weight of $\mathfrak{g}/\mathfrak{k}$. Clearly, a ladder module is multiplicity-free and hence bounded. Moreover, it remains bounded with respect to any $\mathfrak{gl}(2)$ -subalgebra of \mathfrak{k} . Hence any image of $\mathfrak{gl}(2)$ in $\mathfrak{sp}(4)$ or G_2 is bounded.

The only remaining case is $\mathfrak{g} = G_2$, $\mathfrak{k} \simeq \mathfrak{sl}(3)$. To show that \mathfrak{k} is bounded we use Theorem 2.2 with V being the 7-dimensional G_2 -module. Then as a \mathfrak{k} -module V is isomorphic to $V_{\omega_1} \oplus V_{\omega_1}^* \oplus \mathbb{C}$. One can fix a Borel subalgebra $\mathfrak{b} \subset \mathfrak{g}$ so that there exists a \mathfrak{b} -stable one-dimensional subspace $V' \subset V_{\omega_1}^*$. Then $Q_0 \simeq GL(2)$ and

$$(V')^* \otimes (\mathfrak{g} \cdot V' / \mathfrak{k} \cdot V') \simeq \Lambda^2(E) \otimes (E^* \oplus \mathbb{C})$$

where E is the standard $GL(2)$ -module. It is easy to check that it is a spherical Q_0 -module. \square

In the rest of this paper \mathfrak{g} will be of rank 2, and \mathfrak{k} will be isomorphic to $\mathfrak{sl}(2)$. By V_k we denote the $k + 1$ -dimensional \mathfrak{k} -module, and we write $c(M)$ for the \mathfrak{k} -character of any $(\mathfrak{g}, \mathfrak{k})$ -module M of finite type over \mathfrak{k} :

$$c(M) := \sum_{k \geq 0} (\dim M^k) z^k,$$

where $M^k = \text{Hom}_{\mathfrak{k}}(V_k, M)$. By definition, $c(M)$ is a formal power series in z . The *minimal \mathfrak{k} -type* of M is V_t where $t \in \mathbb{Z}_{\geq 0}$ is minimal with $M^t \neq 0$. A $(\mathfrak{g}, \mathfrak{k})$ -module of finite type M is *even* (respectively, *odd*) if $M^t = 0$ for all $t \in 1 + 2\mathbb{Z}$ (resp., $t \in 2\mathbb{Z}$).

Let $\mathbb{C}((z))$ be the algebra of Laurent series and $\mathbb{C}((z))'$ be the span of vectors in $\mathbb{C}((z))$ of the form $z^j + z^{-j-2}$ for $j \in \mathbb{Z}$ ($\mathbb{C}((z))'$ is not a subalgebra). Note that $\mathbb{C}((z))'$ is a complement to the subspace $\mathbb{C}[[z]]$ of $\mathbb{C}((z))$. In what follows we denote by π the projection onto the second summand in the direct sum $\mathbb{C}((z)) = \mathbb{C}((z))' \oplus \mathbb{C}[[z]]$, and we set $z^p \otimes z^q := \sum_{0 \leq k \leq q} z^{p+q-2k}$ for $p \geq q$ and $z^p \otimes z^q := z^q \otimes z^p$ for $p < q$.

Lemma 3.2.

- (a) For any $f(z) \in \mathbb{C}((z))$ and any $j \in \mathbb{Z}$, $\pi(f(z)(z^j + z^{-j})) = \pi(\pi(f(z)(z^j + z^{-j})))$.
- (b) For any $(\mathfrak{k}, \mathfrak{k})$ -module M of finite type over \mathfrak{k}

$$c(M \otimes V_i) = \pi(c(M) \sum_{0 \leq k \leq i} z^{i-2k}),$$

for all $i \in \mathbb{N}$.

Proof.

- (a) It suffices to check that for any $\psi(z) \in \mathbb{C}((z))'$, $\psi(z)(z^j + z^{-j}) \in \mathbb{C}((z))'$, and this is obvious.
- (b) It suffices to check that, for any $s \in \mathbb{Z}_{\geq 0}$

$$\pi(z^s \otimes (\sum_{0 \leq k \leq i} z^{i-2k})) = \sum_{0 \leq k \leq \lfloor \frac{i-s}{2} \rfloor} z^{s+i-2k},$$

which is also obvious.

□

Finally, by $\Gamma_{\mathfrak{k}}$ we denote the functor of \mathfrak{k} -finite vectors:

$$\begin{aligned} \Gamma_{\mathfrak{k}} : \mathfrak{g} - \text{mod} &\rightsquigarrow (\mathfrak{g}, \mathfrak{k}) - \text{mod}, \\ M &\mapsto \{m \in M \mid \dim(U(\mathfrak{k}) \cdot m) < \infty\}. \end{aligned}$$

4. CLASSIFICATION AND \mathfrak{k} -CHARACTERS OF SIMPLE ($\mathfrak{sl}(2) \oplus \mathfrak{sl}(2), \mathfrak{sl}(2)$)-MODULES

Theorem 3.1 singles out the cases when $\mathfrak{k} \simeq \mathfrak{sl}(2)$ is a bounded subalgebra of a rank-2 Lie algebra. The simplest case is when $\mathfrak{g} = \mathfrak{sl}(2) \oplus \mathfrak{sl}(2)$ and $\mathfrak{k} \subset \mathfrak{g}$ is the diagonal subalgebra. Here all simple $(\mathfrak{g}, \mathfrak{k})$ -modules are bounded and are moreover multiplicity-free. This follows, for instance, from the algebraic subquotient theorem, see [Dix], Ch. 9. These $(\mathfrak{g}, \mathfrak{k})$ -modules are historically among the first examples of $(\mathfrak{g}, \mathfrak{k})$ -modules studied. They have been classified already in 1947 by Gelfand and Naimark [GN] and by Bargmann [B], and have been constructed also by Harish-Chandra around the same time, [HC]. A fundamental more modern and much more general reference is the article [BG], where however this explicit example is not written in detail. In the present section we give a quick self-contained description of all simple $(\mathfrak{g}, \mathfrak{k})$ -modules based on the approach of [BG].

Lemma 4.1. *Let $\Omega_1, \Omega_2 \in U(\mathfrak{g})$ be the Casimir elements of the two $\mathfrak{sl}(2)$ -direct summands of \mathfrak{g} , and $\Omega \in U(\mathfrak{k}) \subset U(\mathfrak{g}) = U$ be the Casimir element of \mathfrak{k} . Then Ω_1, Ω_2 and Ω generate $U(\mathfrak{g})^{\mathfrak{k}}$.*

Proof. Straightforward computation. A more general result is proved by F. Knop in [Kn1]. □

Corollary 4.2. *Every simple $(\mathfrak{g}, \mathfrak{k})$ -module is multiplicity-free.*

Denote by $\chi(a, b)$ the central character of the Verma \mathfrak{g} -module with highest weight $(a - 1, b - 1)$, where the notation (c, d) is shorthand for the weight $c\omega_{\text{left}} + d\omega_{\text{right}}$, ω_{left} (respectively, ω_{right}) being the fundamental weight of the first (respectively, second) direct summand of \mathfrak{g} .

Lemma 4.3. *If V_n is the minimal \mathfrak{k} -type of a simple infinite-dimensional $(\mathfrak{g}, \mathfrak{k})$ -module M , then*

$$(2) \quad c(M) = z^n + z^{n+2} + z^{n+4} + \dots \quad .$$

Proof. To prove (2) it suffices to show that V_n, V_{n+2}, V_{n+4} , etc. are precisely all \mathfrak{k} -types of M . The absence of other \mathfrak{k} -types follows from the fact that as a \mathfrak{k} -module \mathfrak{g} is isomorphic to $V_2 \oplus V_2$, hence when acting

by \mathfrak{g} on V_{n+2i} one can only obtain \mathfrak{k} -constituents of $(V_2 \oplus V_2) \otimes V_{n+2i}$, i.e. $V_{n+2(i-1)}$, V_{n+2i} and $V_{n+2(i+1)}$. To show that for each $i > 0$ V_{n+2i} is a \mathfrak{k} -constituent of M , note that if V_{n+2i} were not a constituent of M , then when acting by \mathfrak{g} on $V_{n+2(i-t)}$ for $t \geq 1$ one would not be able to obtain a constituent of the form $V_{n+2(i+r)}$ for $r \geq 1$. Hence M would turn being finite-dimensional, a contradiction. \square

Lemma 4.4. *Let M be a simple $(\mathfrak{g}, \mathfrak{k})$ -module with minimal \mathfrak{k} -type V_0 . Then the central character of M equals $\chi(a, a)$ for some $a \in \mathbb{C}$.*

Proof. Since $\mathfrak{g} \simeq \mathfrak{k} \oplus \mathfrak{k}$, the \mathfrak{g} -module $U \otimes_{U(\mathfrak{k})} V_0$ is isomorphic to $U(\mathfrak{k})$. The latter is endowed with a $U \simeq U(\mathfrak{k}) \otimes U(\mathfrak{k})$ -module structure via left multiplication by elements of $U(\mathfrak{k}) \otimes 1$ and right multiplication by elements of $1 \otimes U(\mathfrak{k})$. Moreover, the action of Ω_1 and Ω_2 coincides on $U(\mathfrak{k})$. Since M is a quotient of the \mathfrak{g} -module $U(\mathfrak{k})$, the action of Ω_1 and Ω_2 coincides on M , hence the Lemma. \square

Lemma 4.5. *Let M be a simple $(\mathfrak{g}, \mathfrak{k})$ -module. Then the central character of M equals $\chi(a, a+n)$ for some $a \in \mathbb{C}$ and some $n \in \mathbb{Z}$. Moreover, the parity of n equals the parity of k where V_k is the minimal \mathfrak{k} -type of M .*

Proof. Let M have central character $\chi(\alpha, \beta)$. Consider the \mathfrak{g} -module $M \otimes (V_0 \boxtimes V_k)$, where the $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{k}$ -module $V_0 \boxtimes V_k$ is endowed with a \mathfrak{g} -module structure via the isomorphism $\mathfrak{g} \simeq \mathfrak{k} \oplus \mathfrak{k}$. Then $\text{Hom}_{\mathfrak{k}}(V_0, M \otimes (V_0 \boxtimes V_k)) \neq 0$, hence a simple subquotient of $M \otimes (V_0 \boxtimes V_k)$ has central character $\chi(a, a)$ for some a . On the other hand, the central characters of all simple subquotients of $M \otimes (V_0 \boxtimes V_k)$ are of the form $\chi(\alpha, \beta - n)$ for n running over the set of weights of V_k . Therefore $\alpha = a$, $\beta - n = a$, i.e. the Lemma follows. \square

Lemma 4.6. *For any central character χ , up to isomorphism there is at most one infinite dimensional simple $(\mathfrak{g}, \mathfrak{k})$ -module with this central character.*

Proof. Let M', M'' be two simple $(\mathfrak{g}, \mathfrak{k})$ -modules with central character χ . Then, by Lemma 4.3, for some m $\text{Hom}_{\mathfrak{k}}(V_m, M') = \text{Hom}_{\mathfrak{k}}(V_m, M'') = \mathbb{C}$. Therefore M' and M'' are isomorphic to simple quotients of the \mathfrak{g} -module $U \otimes_{Z_U U(\mathfrak{k})} V_m$, where Z_U acts on V_m via the central character χ . The fact that $U^{\mathfrak{k}} \subset Z_U U(\mathfrak{k})$ (Lemma 4.1) implies that $\text{Hom}_{\mathfrak{k}}(V_m, U \otimes_{Z_U U(\mathfrak{k})} V_m) = \mathbb{C}$ for every $m \geq 0$. Hence $U \otimes_{Z_U U(\mathfrak{k})} V_m$ has a unique proper maximal submodule, and in this way also a unique simple quotient. Therefore $M' \simeq M''$. \square

In the rest of this section we only consider central characters of the form $\chi(a, a - n)$ for $n \in \mathbb{Z}_{\geq 0}$. If $a \in \mathbb{Z}$, we assume in addition that

$a \geq 0$ and $a - n \leq 0$. By M_c denote the Verma module over \mathfrak{k} with highest weight $c - 1$. Note that for $a, a - n$ as above, $\text{Hom}_{\mathbb{C}}(M_a, M_{a-n})$ is a \mathfrak{g} -module with central character $\chi(a, a - n)$. Define

$$W_{a,a-n} := \Gamma_{\mathfrak{k}}(\text{Hom}_{\mathbb{C}}(M_a, M_{a-n})).$$

Theorem 4.7.

- (a) Fix $a \in \mathbb{C} \setminus \mathbb{Z}_{<0}$ and $n \in \mathbb{Z}_{\geq 0}$ such that $a - n \leq 0$ for integer a . The \mathfrak{g} -module $W_{a,a-n}$ is the unique (up to isomorphism) simple infinite-dimensional $(\mathfrak{g}, \mathfrak{k})$ -module with central character $\chi(a, a - n)$.
- (b) $c(W_{a,a-n}) = z^n + z^{n+2} + z^{n+4} + \dots$.

Proof. Note that to compute the \mathfrak{k} -character of $\Gamma_{\mathfrak{k}}(\text{Hom}_{\mathbb{C}}(M_a, M_{a-n}))$ it suffices to compute $\text{Hom}_{\mathfrak{k}}(V_m, \text{Hom}_{\mathbb{C}}(M_a, M_{a-n}))$ for all $m \in \mathbb{Z}_{\geq 0}$. However,

$$\text{Hom}_{\mathfrak{k}}(V_m, \text{Hom}_{\mathbb{C}}(M_a, M_{a-n})) = \text{Hom}_{\mathfrak{k}}(M_a, M_{a-n} \otimes V_m^*),$$

and

$$\text{Hom}_{\mathfrak{k}}(M_a, M_{a-n} \otimes V_m^*) = \begin{cases} \mathbb{C} & \text{for } m - n \in 2\mathbb{Z}_{\geq 0} \\ 0 & \text{otherwise} \end{cases} .$$

Hence

$$c(W_{a,a-n}) = z^n + z^{n+2} + z^{n+4} + \dots .$$

The simplicity of $W_{a,a-n}$ follows from the observation that if reducible, $W_{a,a-n}$ would have a finite-dimensional subquotient, but there is no finite-dimensional \mathfrak{g} -module with central character $\chi(a, a - n)$ for $a \in \mathbb{C} \setminus \mathbb{Z}$ or $a = 0$. If $a \in \mathbb{Z}$, the finite-dimensional \mathfrak{g} -module with central character $\chi(a, a - n)$ is isomorphic to $V_{a-1} \boxtimes V_{n-a-1}$ whose \mathfrak{k} -character is $z^{n-2} + z^{n-4} + \dots + z^{|n-2a-2|}$, and hence it can not be a subquotient of $W_{a,a-n}$. \square

5. CLASSIFICATION AND \mathfrak{k} -CHARACTERS OF SIMPLE BOUNDED ($\mathfrak{sl}(3), \mathfrak{sl}(2)$)-MODULES

Throughout this section $\mathfrak{g} = \mathfrak{sl}(3)$ and $\mathfrak{k} \simeq \mathfrak{sl}(2) \subset \mathfrak{g}$.

5.1. The root case. In this subsection we fix a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$ and simple roots $\alpha_1, \alpha_2 \in \mathfrak{h}^*$ which define a Borel subalgebra $\mathfrak{b}^+ \subset \mathfrak{g}$. We also fix \mathfrak{k} to be the $\mathfrak{sl}(2)$ -subalgebra generated by the root spaces $\mathfrak{g}^{\pm\alpha_1}$. There are two parabolic subalgebras containing \mathfrak{k} and \mathfrak{h} : $\mathfrak{p}^+ := (\mathfrak{h} + \mathfrak{k}) \oplus \mathfrak{g}^{\alpha_2} \oplus \mathfrak{g}^{\alpha_1+\alpha_2}$, $\mathfrak{p}^- := (\mathfrak{h} + \mathfrak{k}) \oplus \mathfrak{g}^{-\alpha_2} \oplus \mathfrak{g}^{-\alpha_1-\alpha_2}$. Note that $\mathfrak{b}^+ \subset \mathfrak{p}^+$ and define \mathfrak{b}^- to be the Borel subalgebra with simple roots $\alpha_1, -\alpha_1 - \alpha_2$. Then $\mathfrak{b}^- \subset \mathfrak{p}^-$. In addition, we fix generators $h_i \in [\mathfrak{g}^{\alpha_i}, \mathfrak{g}^{-\alpha_i}]$ and denote by ω_i , for $i = 1, 2$, the corresponding dual basis of \mathfrak{h}^* . Then $\rho_{\mathfrak{b}^+} = \omega_1 + \omega_2$, $\rho_{\mathfrak{b}^-} = \omega_1 - 2\omega_2$.

Lemma 5.1. *Let M be a simple bounded infinite-dimensional $(\mathfrak{g}, \mathfrak{k})$ -module. Then $\mathfrak{g}[M] = \mathfrak{p}^\pm$.*

Proof. Since $\mathfrak{h} \subset \mathfrak{g}^\mathfrak{k} \oplus \mathfrak{k}$, we have $\mathfrak{h} \subset \mathfrak{g}[M]$. Put $M_0 := \{m \in M \mid \mathfrak{g}^{\alpha_1} \cdot m = 0\}$ and choose generators x and y of the respective root spaces $\mathfrak{g}^{-\alpha_2}$ and $\mathfrak{g}^{\alpha_1 + \alpha_2}$. A straightforward computation shows that for any $i, j \in \mathbb{Z}_{\geq 0}$, $(x^i y^j) \cdot v \in M_0$ if v is any non-zero vector in M_0 such that $h_1 \cdot v = \nu(h_1)v$ for some $\nu \in (\mathfrak{h} \cap \mathfrak{k})^*$. Therefore the assumption that $x, y \notin \mathfrak{g}[M]$ implies that the multiplicity of $V_{\nu+i+j}$ is at least $i+j$, which contradicts the boundedness of M . Hence $\mathfrak{g}^{-\alpha_2} \in \mathfrak{g}[M]$ or $\mathfrak{g}^{\alpha_1 + \alpha_2} \in \mathfrak{g}[M]$, and consequently $\mathfrak{g}[M] = \mathfrak{p}^\pm$. \square

Let $F_{a,b}^\pm$ be the simple finite-dimensional \mathfrak{p}^\pm -module with \mathfrak{b}^\pm -highest weight $a\omega_1 + b\omega_2$. Define $L_{a,b}^\pm$ as the unique simple quotient of $U(\mathfrak{g}) \otimes_{U(\mathfrak{p}^\pm)} F_{a,b}^\pm$. Then $L_{a,b}^\pm$ are bounded $(\mathfrak{g}, \mathfrak{k})$ -modules, and the existence of an isomorphism $L_{a,b}^\pm \simeq L_{a',b'}^\mp$ implies $\dim L_{a,b}^\pm < \infty$.

Theorem 5.2. *Let, as above, $\mathfrak{k} \simeq \mathfrak{sl}(2)$ be a root subalgebra of $\mathfrak{g} = \mathfrak{sl}(3)$.*

(a) *Any infinite-dimensional bounded $(\mathfrak{g}, \mathfrak{k})$ -module is isomorphic either to $L_{a,b}^+$ for $a \in \mathbb{Z}_{\geq 0}$, $b \in \mathbb{C} \setminus \mathbb{Z}_{\geq 0}$ or to $L_{a,b}^-$ for $a \in \mathbb{Z}_{\geq 0}$, $-a - b \in \mathbb{C} \setminus \mathbb{Z}_{\geq 0}$.*

(b)

$$(3) \quad c(L_{a,b}^\pm) = 1 + 2z + \cdots + az^{a-1} + (a+1)(z^a + z^{a+1} + \cdots)$$

for all $a \geq 0$ and for those b which do not satisfy the conditions $-b \in \mathbb{Z}_{\geq 2}$, $a+b \in \mathbb{Z}_{\geq -1}$ for $L_{a,b}^+$, and respectively the conditions $a+b \in \mathbb{Z}_{\geq 2}$, $-b \in \mathbb{Z}_{\geq -1}$ for $L_{a,b}^-$.

(c) *If $-b \in \mathbb{Z}_{\geq 2}$, $a+b \in \mathbb{Z}_{\geq -1}$, then*

$$(4) \quad c(L_{a,b}^+) = z^{-b-1} + 2z^{-b} + \cdots + (a+b+1)z^{a-1} + (a+b+2)(z^a + z^{a+1} + \cdots),$$

and if $a+b \in \mathbb{Z}_{\geq 2}$, $-b \in \mathbb{Z}_{\geq -1}$, then

$$(5) \quad c(L_{a,b}^-) = z^{a+b-1} + 2z^{a+b} + \cdots + (1-b)z^{a-1} + (2-b)(z^a + z^{a+1} + \cdots).$$

Proof. Let M be a simple infinite-dimensional bounded $(\mathfrak{g}, \mathfrak{k})$ -module. Then, by Lemma 5.1, $\mathfrak{g}[M] = \mathfrak{p}^\pm$. If $\mathfrak{g}[M] = \mathfrak{p}^+$, let M^+ be a simple finite-dimensional \mathfrak{p}^+ -submodule of M . Then $M^+ \simeq F_{a,b}^+$ for some $a \in \mathbb{Z}_{\geq 0}$ and some $b \in \mathbb{C}$, and there is an obvious surjection of \mathfrak{g} -modules $U(\mathfrak{g}) \otimes_{U(\mathfrak{p}^+)} F_{a,b}^+ \rightarrow M$. Hence M is isomorphic to the unique simple quotient $L_{a,b}^+$ of $U(\mathfrak{g}) \otimes_{U(\mathfrak{p}^+)} F_{a,b}^+$. However, $L_{a,b}^+$ is finite-dimensional iff $b \in \mathbb{Z}_{\geq 0}$, therefore (a) follows for the case when $\mathfrak{g}[M] = \mathfrak{p}^+$. The case $\mathfrak{g}[M] = \mathfrak{p}^-$ is obtained by replacing b with $-a - b$ which corresponds to

the replacement of the simple root α_2 of \mathfrak{b}^+ by the simple root $-\alpha_1 - \alpha_2$ of \mathfrak{b}^- .

Statements (b) and (c) follow from a non-difficult reducibility analysis for the induced module $U(\mathfrak{g}) \otimes_{U(\mathfrak{p}^\pm)} F_{a,b}^\pm$. Note first of all that $\text{ch}_{\mathfrak{k}}(U(\mathfrak{g}) \otimes_{U(\mathfrak{p}^\pm)} F_{a,b}^\pm)$ is always given by the right-hand side of (3). Indeed as \mathfrak{k} -modules $\mathfrak{g}/\mathfrak{p}^\pm$ and $F_{a,b}^\pm$ are isomorphic respectively to V_1 and V_a , therefore

$$c(U(\mathfrak{g}) \otimes_{U(\mathfrak{p}^\pm)} F_{a,b}^\pm) = c(S(V_1) \otimes V_a).$$

A straightforward computation shows that $c(S(V_1) \otimes V_a)$ is nothing but the right hand side of (3).

We claim now that $U(\mathfrak{g}) \otimes_{U(\mathfrak{p}^\pm)} F_{a,b}^\pm$ is irreducible precisely when b does not satisfy the respective conditions stated in (b). Consider first the case of \mathfrak{p}^+ . Then $U(\mathfrak{g}) \otimes_{U(\mathfrak{p}^+)} F_{a,b}^+$ is irreducible if and only if there exists $w \in W \setminus W_{\mathfrak{k}}$ such that

$$(6) \quad (w((a+1)\omega_1 + (b+1)\omega_2) - (\omega_1 + \omega_2))(h_1) \in \mathbb{Z}_{\geq 0}$$

and

$$(7) \quad (w((a+1)\omega_1 + (b+1)\omega_2) - (\omega_1 + \omega_2)) = a\omega_1 + b\omega_2 - m_1\alpha_1 - m_2\alpha_2$$

for some $m_1, m_2 \in \mathbb{Z}_{\geq 0}$. The only non \mathfrak{b}^+ -dominant solution of (6) and (7) is $w = w_{\alpha_1 + \alpha_2}$ and $-b \in \mathbb{Z}_{\geq 2}$, $a + b \in \mathbb{Z}_{\geq -1}$. Moreover, in the latter case $L_{a,b}^+ \simeq (U(\mathfrak{g}) \otimes_{U(\mathfrak{p}^+)} F_{a,b}^+) / L_{-b-2, -a-2}^+$, where $c(L_{-b-2, -a-2}^+)$ is given by the right hand side of (3) with a replaced by $-b-2$. An immediate computation shows that $c(L_{a,b}^+)$ is given in this case by the right hand side of (4), therefore (b) and (c) are proved for the case of \mathfrak{p}^+ . The case of \mathfrak{p}^- is obtained by interchanging the parameter b in (4) with $-a-b$. \square

Corollary 5.3. *Let \mathfrak{g} and \mathfrak{k} be as above.*

- (a) *The minimal \mathfrak{k} -type of a simple bounded infinite-dimensional $(\mathfrak{g}, \mathfrak{k})$ -module can be arbitrary. The multiplicity of the minimal \mathfrak{k} -type is always 1.*
- (b) *The following is a complete list of multiplicity-free simple infinite-dimensional $(\mathfrak{g}, \mathfrak{k})$ -modules:*
 - $L_{0,b}^+$ for $b \in \mathbb{C} \setminus \mathbb{Z}_{\geq 0}$,
 - $L_{0,b}^-$ for $-b \in \mathbb{C} \setminus \mathbb{Z}_{\geq 0}$,
 - $L_{a,b}^+$ for $a + b = -1$, $-b \in \mathbb{Z}_{\geq 2}$,
 - $L_{a,b}^-$ for $b = 1$, $a + b \in \mathbb{Z}_{\geq 2}$.

5.2. The principal case. Let now \mathfrak{k} be a principal $\mathfrak{sl}(2)$ -subalgebra of $\mathfrak{g} = \mathfrak{sl}(3)$. The pair $(\mathfrak{g}, \mathfrak{k})$ is well known to be symmetric and the simple $(\mathfrak{g}, \mathfrak{k})$ -modules have been studied extensively, see for instance [Fo] and [Sp]. In principle one should be able to identify all simple bounded modules in the known classification of simple Harish-Chandra modules. However, we propose an alternative approach which leads directly to all bounded simple $(\mathfrak{g}, \mathfrak{k})$ -modules and their \mathfrak{k} -characters. This is the first case in which the richness of the theory of bounded (generalized) Harish-Chandra modules becomes apparent.

We keep the notations $\mathfrak{h}, \mathfrak{b}^+, \alpha_1, \alpha_2$ from Subsection 5.1. By $L_{a,b}$ we denote the simple \mathfrak{g} -module with \mathfrak{b}^+ -highest weight $(a-1)\omega_1 + (b-1)\omega_2$, by $V_{p,q}$ we denote the simple finite-dimensional $\mathfrak{g} = \mathfrak{sl}(3)$ -module with \mathfrak{b}^+ -highest weight $p\omega_1 + q\omega_2$ ($p, q \in \mathbb{Z}_{\geq 0}$), and $\chi(a, b)$ stands for the central character of $L_{a,b}$. By A we denote the Weyl algebra in the indeterminates t, x, y .

We first describe the primitive ideals of all simple bounded $(\mathfrak{g}, \mathfrak{k})$ -modules. Let $\mathrm{GKdim}M$ denote the Gelfand-Kirillov dimension of a \mathfrak{g} -module M and X_M denote the associated variety of M .

Lemma 5.4. *Let M be an infinite-dimensional bounded simple $(\mathfrak{g}, \mathfrak{k})$ -module. Then $\mathrm{Ann}M = \mathrm{Ann}L_{a,b}$, where $\dim L_{a,b} = \infty$, $a \in \mathbb{Z}_{>0}$, $b \in \mathbb{Z}_{>0}$ or $a + b \in \mathbb{Z}_{>0}$.*

Proof. By Duflo's Theorem $\mathrm{Ann}M = \mathrm{Ann}L_{a,b}$ for some a, b . By Theorem 4.4 in [PS2], $\mathrm{GKdim}M \leq 2$. Since $\mathrm{GKdim}M \geq \frac{1}{2}\dim X_M$ and $\mathrm{GKdim}L_{a,b} = \frac{1}{2}\dim X_M$, we have $\mathrm{GKdim}L_{a,b} \leq 2$. A straightforward computation shows that this latter condition is equivalent to the condition on (a, b) in the statement of the Lemma. \square

Let $\mathfrak{B}_{\mathfrak{k}}^{\chi}$ be the category of bounded $(\mathfrak{g}, \mathfrak{k})$ -modules which afford the central character χ , see [PS2], Section 4.

Corollary 5.5. *If $\mathfrak{B}_{\mathfrak{k}}^{\chi}$ is not empty, then $\chi = \chi(u + 1 - n, n + 1)$ for some $n \in \mathbb{Z}_{\geq 0}$, where $u \in \mathbb{C} \setminus \mathbb{Z}_{< n-1}$ or $u = -2$.*

Note that the natural embedding of $\mathfrak{gl}(3)$ into A maps the center of $\mathfrak{gl}(3)$ to the line $\mathbb{C}\mathbf{E}$ for $\mathbf{E} := t\partial_t + x\partial_x + y\partial_y$, and that the adjoint action of the central element \mathbf{E} on A defines a \mathbb{Z} -grading $A := \bigoplus_{i \in \mathbb{Z}} A_i$. Let $u \in \mathbb{C}$. Define the (associative) algebra D^u as the quotient of A_0 by the ideal generated by $\mathbf{E} - u$. The embedding of $\mathfrak{g} \rightarrow A_0$ induces a surjective homomorphism $\gamma_u : U(\mathfrak{g}) \rightarrow D^u$. It is not difficult to show that if $u \in \mathbb{Z}$, D^u is isomorphic to the algebra of globally defined differential endomorphisms of the line bundle $\mathcal{O}_{\mathbb{P}^2}(u)$ (\mathbb{P}^2 being the projective space with homogeneous coordinates (x, y, z)).

Lemma 5.6. *Consider D^u with its adjoint \mathfrak{g} -module structure. Then*

$$D^u \simeq \bigoplus_{m \geq 0} V_{m\rho}.$$

Proof. Let $\mathbb{C} = A^0 \subset A^1 \subset \dots \subset A$ denote the standard filtration of A . A direct computation shows that as a \mathfrak{g} -module A_0^m/A_0^{m-1} is isomorphic to

$$V_{m,0} \otimes V_{0,m} = \bigoplus_{k=0}^m V_{k\rho}.$$

After factorization by $\mathbf{E} - u$, one obtains

$$(D^u)^m / (D^u)^{m-1} \simeq V_{m\rho}.$$

□

It is not difficult to see that the restriction of γ_u to $U(\mathfrak{k})$ is injective. Slightly abusing notation we identify $U(\mathfrak{k})$ with its image in D^u . We will use the following expression for the standard basis E, H, F of \mathfrak{k} :

$$(8) \quad E = t\partial_x + x\partial_y, H = 2t\partial_t - 2y\partial_y, F = 2x\partial_t + 2y\partial_x.$$

Lemma 5.7. *The centralizer of \mathfrak{k} in D^u coincides with the center of $U(\mathfrak{k}) \subset D^u$.*

Proof. As $V_{m\rho}^{\mathfrak{k}} = 0$ for odd m and $V_{m\rho}^{\mathfrak{k}} = \mathbb{C}$ for even m it is clear that the centralizer of \mathfrak{k} in D^u is generated by the quadratic Casimir element $\Omega \in V_{2\rho}^{\mathfrak{k}}$. □

Corollary 5.8. *Every simple (D^u, \mathfrak{k}) -module is multiplicity-free. For any non-negative m , there exists at most one (up to isomorphism) simple (D^u, \mathfrak{k}) -module M with $\text{Hom}_{\mathfrak{k}}(V_m, M) \neq 0$.*

Proof. It is well known that if M is a simple $(\mathfrak{g}, \mathfrak{k})$ -module, then $M^V = \text{Hom}_{\mathfrak{k}}(V, M)$ is a simple $U(\mathfrak{g})^{\mathfrak{k}}$ -module for every \mathfrak{k} -type V , see for instance Lemma 3.3 in [PS2]. Therefore Lemma 5.7 implies the first statement. The proof of the second statement is very similar to the proof of Lemma 4.6. □

We now introduce the functors

$$\begin{aligned} \text{Ind} : D^u - \text{mod} &\hookrightarrow A - \text{mod} \\ M &\mapsto A \otimes_{A_0} M, \\ \text{Res}_u : A - \text{mod} &\hookrightarrow D^u - \text{mod} \\ M &\mapsto D^u \otimes_{A_0} M. \end{aligned}$$

Obviously, $\text{Res}_u \circ \text{Ind} = \text{id}_{D^u - \text{mod}}$.

Lemma 5.9.

$$\ker \gamma_u = \begin{cases} \text{Ann}L_{u+1,1} = \text{Ann}L_{-u-1,u+2} = \text{Ann}L_{1,-u-2} & \text{for } u \notin \mathbb{Z} \\ \text{Ann}L_{-u-1,u+2} = \text{Ann}L_{1,-u-2} & \text{for } u \in \mathbb{Z}_{\geq -1} \\ \text{Ann}L_{u+1,1} = \text{Ann}L_{-u-1,u+2} & \text{for } u \in \mathbb{Z}_{\leq -2} \end{cases} .$$

Proof. First we prove that $\ker \gamma_u \subset \text{Ann}L_{a,b}$ with a, b as in the statement. Note that $\text{Res}_u(t^u \mathbb{C}[t^{\pm 1}, x, y])$ contains a submodule generated by t^u isomorphic to $L_{u+1,1}$, $\text{Res}_u(x^u \mathbb{C}[t^{\pm 1}, x^{\pm 1}, y]) / \text{Res}_u(x^u \mathbb{C}[t, x^{\pm 1}, y])$ contains a submodule with highest vector $t^{-1}x^{u+1}$ isomorphic to $L_{-u-1,u+2}$ and $\text{Res}_u(y^u \mathbb{C}[t^{\pm 1}, x^{\pm 1}, y^{\pm 1}]) / (\text{Res}_u(y^u \mathbb{C}[t^{\pm 1}, x, y^{\pm 1}]) + \text{Res}_u(y^u \mathbb{C}[t, x^{\pm 1}, y^{\pm 1}]))$ contains a submodule with highest vector $t^{-1}x^{-1}y^{u+2}$ isomorphic to $L_{1,-u-2}$. Hence $\ker \gamma_u \subset \text{Ann}L_{a,b}$. Next we see from Lemma 5.6 that all proper two-sided ideals of D^u have finite codimension. Thus, $\gamma_u(\text{Ann}L_{a,b})$ is either 0 or has finite codimension in D^u . The latter is impossible because $L_{a,b}$ is infinite-dimensional. Hence $\ker \gamma_u = \text{Ann}L_{a,b}$. \square

Since the eigenvalues of ad_H in $U(\mathfrak{g})$ are all even, every simple $(\mathfrak{g}, \mathfrak{k})$ -module is either odd or even.

As follows from Lemma 5.9, all simple bounded $(\mathfrak{g}, \mathfrak{k})$ -modules with central character $\chi(u+1, u)$ are (D^u, \mathfrak{k}) -modules. This allows us to first classify the simple (D^u, \mathfrak{k}) -modules and then use translation functors to classify the bounded simple modules with arbitrary possible central character, see Corollary 5.5.

Note that the functor Ind maps (D^u, \mathfrak{k}) -mod into $(\widetilde{A}, \mathfrak{k})$ -mod, the latter being defined as the full subcategory of A -mod consisting of \mathfrak{k} -locally finite A -modules with semisimple action of \mathbf{E} .

Lemma 5.10. *For any simple (D^u, \mathfrak{k}) -module M there exists a simple $(\widetilde{A}, \mathfrak{k})$ -module \hat{M} with $\text{Res}_u(\hat{M}) \simeq M$.*

Proof. Let N be a maximal proper A -submodule of $\text{Ind}(M)$. Then $\text{Res}_u(N) \not\simeq M$ as M generates $\text{Ind}(M)$. Therefore $\text{Res}_u(N) = 0$ and one defines \hat{M} as $\text{Ind}(M)/N$. \square

Set $f := x^2 - 2ty$, $\Delta := \partial_x^2 - 2\partial_y \partial_t$ and note that $f, \Delta \in A^\mathfrak{k}$. For every fixed $p \in \mathbb{C}$, we put $R^p := f^p \mathbb{C}[t, x, y, f^{-1}]$. Then clearly R^p is an (A, \mathfrak{k}) -module and $\text{Res}_u(R^p) = 0$ if $u - 2p \notin \mathbb{Z}$. Otherwise,

$$\text{Res}_u(R^p) = \begin{cases} \mathbb{C}f^{\frac{u}{2}} \oplus f^{\frac{u-2}{2}} \mathcal{H}_2 \oplus f^{\frac{u-4}{2}} \mathcal{H}_4 \oplus \dots & \text{for } u - 2p \in 2\mathbb{Z} \\ \mathbb{C}f^{\frac{u-1}{2}} \mathcal{H}_1 \oplus f^{\frac{u-3}{2}} \mathcal{H}_3 \oplus f^{\frac{u-5}{2}} \mathcal{H}_5 \oplus \dots & \text{for } u - 2p \in 2\mathbb{Z} + 1, \end{cases}$$

where \mathcal{H}_n denotes the space of homogeneous polynomials of degree n in $\mathbb{C}[t, x, y]$ annihilated by Δ (as a \mathfrak{k} -module \mathcal{H}_n is isomorphic to V_{2n}).

Lemma 5.11.

- (a) For $u \notin \mathbb{Z}$ and for $u = -1, -2$, $\text{Res}_u(R^{\frac{u}{2}})$ and $\text{Res}_u(R^{\frac{u+1}{2}})$ are simple D^u -modules.
- (b) For $u \in 2\mathbb{Z}_{\geq 0}$, $\text{Res}_u(R^{\frac{u+1}{2}})$ is a simple D^u -module and there is an exact sequence

$$(10) \quad 0 \rightarrow V_{u,0} \rightarrow \text{Res}_u(R^{\frac{u}{2}}) \rightarrow I_{u,0}^+ \rightarrow 0$$

for some simple D^u -module $I_{u,0}^+$.

- (c) For $u \in 1 + 2\mathbb{Z}_{\geq 0}$, $\text{Res}_u(R^{\frac{u}{2}})$ is a simple D^u -module and there is an exact sequence

$$0 \rightarrow V_{u,0} \rightarrow \text{Res}_u(R^{\frac{u+1}{2}}) \rightarrow I_{u,0}^- \rightarrow 0$$

for some simple D^u -module $I_{u,0}^-$.

- (d) For $u \in 2\mathbb{Z}_{\leq -2}$, $\text{Res}_u(R^{\frac{u}{2}})$ is a simple D^u -module and there is an exact sequence

$$0 \rightarrow I_{u,0}^- \rightarrow \text{Res}_u(R^{\frac{u+1}{2}}) \rightarrow V_{0,-3-u} \rightarrow 0$$

for some simple D^u -module $I_{u,0}^-$.

- (e) For $u \in 1 + 2\mathbb{Z}_{\leq -1}$, $\text{Res}_u(R^{\frac{u+1}{2}})$ is a simple D^u -module and there is an exact sequence

$$0 \rightarrow I_{u,0}^+ \rightarrow \text{Res}_u(R^{\frac{u}{2}}) \rightarrow V_{0,-3-u} \rightarrow 0$$

for some simple D^u -module $I_{u,0}^+$.

Proof. The isomorphism (9) yields

$$(11) \quad c(\text{Res}_u(R^{\frac{u}{2}})) = 1 + z^4 + z^8 + \dots, \quad c(\text{Res}_u(R^{\frac{u+1}{2}})) = z^2 + z^6 + z^{10} + \dots$$

Thus, if $\text{Res}_u(R^{\frac{u}{2}})$ (respectively $\text{Res}_u(R^{\frac{u+1}{2}})$) is not simple it has a unique simple finite-dimensional submodule or a unique simple finite-dimensional quotient. By Lemma 5.9 the latter can happen only if $u \in \mathbb{Z}_{\geq 0}$ or $u \in \mathbb{Z}_{\leq -3}$. Hence (a).

Let $u \in 2\mathbb{Z}_{\geq 0}$. Then $\text{Res}_u(R^{\frac{u}{2}})$ contains $\text{Res}_u(\mathbb{C}[t, x, y]) \simeq V_{u,0}$ as a finite-dimensional simple submodule, hence (10). The \mathfrak{g} -module $\text{Res}_u(R^{\frac{u+1}{2}})$ has the same central character as $\text{Res}_u(R^{\frac{u}{2}})$ and, since $V_{n,0}$ is not a subquotient of $\text{Res}_u(R^{\frac{u+1}{2}})$ by (11), $\text{Res}_u(R^{\frac{u+1}{2}})$ is a simple D^u -module. Hence (b).

As $\Delta(f^{-\frac{1}{2}}) = 0$, $f^{-\frac{1}{2}}$ generates a proper A -submodule $M \subset f^{\frac{1}{2}}\mathbb{C}[t, x, y, f^{-1}]$. A direct computation shows that $\dim \text{Res}_u(M) = \infty$ for any $u \in 1 + 2\mathbb{Z}_{\geq -2}$. Furthermore, the only finite-dimensional module, whose

central character coincides with that of D^u is $V_{0,-3-u}$. Therefore one necessarily has

$$0 \rightarrow I_{u,0}^+ \rightarrow \text{Res}_u(R^{\frac{u}{2}}) \rightarrow V_{0,-3-u} \rightarrow 0$$

where $I_{u,0}^+ := \text{Res}_u(M)$. $\text{Res}_u(R^{\frac{u+1}{2}})$ is simple by the same reason as in (b). Hence (e).

(c) and (d) are similar to (b) and (e). \square

For any $u \in \mathbb{C}$ we define now $I_{u,0}^+$ (respectively, $I_{u,0}^-$) as the unique simple infinite-dimensional constituent of $\text{Res}_u(R^{\frac{u}{2}})$ (resp., $\text{Res}_u(R^{\frac{u+1}{2}})$).

Corollary 5.12. *Every simple even infinite-dimensional (D^u, \mathfrak{k}) -module is isomorphic to $I_{u,0}^\pm$.*

Proof. For every fixed u and any sufficiently large $m \in 2\mathbb{Z}_{\geq 0}$ (such that V_m is not a \mathfrak{k} -type of $V_{u,0}$ or $V_{0,-3-u}$ for $u \in \mathbb{Z}$), Lemma 5.11 implies $\text{Hom}_{\mathfrak{k}}(V_m, I_u^\pm) \neq 0$. The statement follows now from Corollary 5.8. \square

Lemma 5.13. *If $u \notin \frac{1}{2} + \mathbb{Z}$, then every (D^u, \mathfrak{k}) -module is even.*

Proof. Assume that M is an odd simple (D^u, \mathfrak{k}) -module and $u \notin \frac{1}{2} + \mathbb{Z}$. Let \hat{M} be as in Lemma 5.10, A_f denote the localization of A in f , $\hat{M}_f = A_f \otimes_A \hat{M}$. First, we claim that if $u \notin \frac{1}{2} + \mathbb{Z}$, then $\hat{M}_f \neq 0$. Indeed, $\hat{M}_f = 0$ implies that f acts locally nilpotently on \hat{M} . Then $M^0 := \ker f$ is a \mathfrak{k} -submodule of \hat{M} and a straightforward calculation using (8) shows $\Omega_{|M^0} = 2(\mathbf{E} + 3)(\mathbf{E} + 2)|_{M^0}$. Thus $\text{Hom}_{\mathfrak{k}}(V_m, M^0) \neq 0$ only if $2(d+3)(d+2) = \frac{m^2}{2} + m$ or equivalently $(d + \frac{5}{2})^2 = (\frac{m+1}{2})^2$, where d is the eigenvalue of \mathbf{E} on M^0 . Since $d \in u + \mathbb{Z}$, $u \notin \frac{1}{2} + \mathbb{Z}$ implies $M^0 = 0$.

Our next observation is that \hat{M}_f is an odd (A, \mathfrak{k}) -module and that t does not act locally nilpotently on \hat{M}_f . Indeed, if t acts locally nilpotently, by \mathfrak{k} -invariance x and y act locally nilpotently, and therefore f acts locally nilpotently. Contradiction. Therefore \hat{M}_f is a submodule of its localization in t , $\hat{M}_{f,t}$. Furthermore, for some odd m there exists a non-zero vector $v \in \hat{M}_{f,t}$ such that $H \cdot v = mv$, $E \cdot v = 0$ and $\mathbf{E} \cdot v = uv$. The expressions for E, H and \mathbf{E} imply

$$\partial_t v = \frac{-(u + m/2)ty + mx^2/2}{tf}v, \partial_x v = \frac{(u - m/2)x}{f}v, \partial_y v = \frac{(m/2 - u)t}{f}v.$$

Thus, every vector in $\hat{M}_{f,t}$ can be obtained from v by applying elements of $\mathbb{C}[t^{\pm 1}, x, y, f^{-1}]$, i.e. $\hat{M}_{f,t} = \mathbb{C}[t^{\pm 1}, x, y, f^{-1}]v$. It is not difficult to see that $v = t^{\frac{m}{2}} f^{\frac{2u-m}{4}}$ satisfies the above relations. The $A_{f,t}$ -module

$\mathbb{C}[t^{\pm 1}, x, y, f^{-1}]v$ is simple and free over $\mathbb{C}[t^{\pm 1}, x, y, f^{-1}]$. Hence $\hat{M}_{f,t} \simeq \mathbb{C}[t^{\pm 1}, x, y, f^{-1}]v$ and it is obvious that $\hat{M}_{f,t}$ has no non-zero \mathfrak{k} -finite vectors. As we pointed out above, $\hat{M}_f \subset \hat{M}_{f,t}$. Therefore $\hat{M}_f = 0$. \square

We now turn to odd simple (D^u, \mathfrak{k}) -modules.

Lemma 5.14. *Let $u \in \frac{1}{2} + \mathbb{Z}$. Up to isomorphism, there exists exactly one odd simple (D^u, \mathfrak{k}) -module $J_{u,0}$. Moreover,*

$$(12) \quad c(J_{u,0}) = \begin{cases} z^{2-2u} + z^{6-2u} + z^{10-2u} + \dots & \text{for } u < 0 \\ z^{4+2u} + z^{8+2u} + z^{12+2u} + \dots & \text{for } u > 0 \end{cases} .$$

Proof. Let $P \subset G = SL(3)$ be the maximal parabolic subgroup whose Lie algebra \mathfrak{p} equals $\mathfrak{b} \oplus \mathfrak{g}^{-\alpha_1}$, $K \subset G$ be the algebraic subgroup with Lie algebra \mathfrak{k} , and Z be the closed K -orbit on $G/P \simeq \mathbb{P}^2$. Then $Z \simeq \mathbb{P}^1$ and the embedding $i : Z \rightarrow \mathbb{P}^2$ is a Veronese embedding of degree 2. It is not difficult to verify that the relative tangent bundle \mathcal{T}_P of the projection $p : G/B \rightarrow G/P$ is a $\mathcal{O}_{G/B}$ -submodule of the twisted sheaf of differential operators $\mathcal{D}_{G/B}^{(u+1)\omega_1+\omega_2}$ (the definition of $\mathcal{D}_{G/B}^\zeta$ see in [PS2], Section 5). Furthermore, the direct image $p_*(\mathcal{D}_{G/B}^{(u+1)\omega_1+\omega_2}/\mathcal{I}_P)$, where \mathcal{I}_P is the left ideal in $\mathcal{D}_{G/B}^{(u+1)\omega_1+\omega_2}$ generated by \mathcal{T}_P , is a well-defined twisted sheaf of differential operators on G/P . We denote this sheaf by $\mathcal{D}_{G/P}^{(u+1)\omega_1+\omega_2}$.

Our next observation is that, similarly to the equivalence of categories i_\star discussed in Section 5 of [PS2], Kashiwara's theorem yields an equivalence of categories

$$i_\star^u : \mathcal{O}_Z(2u) \otimes_{\mathcal{O}_{G/P}} \mathcal{D}_{G/P} \otimes_{\mathcal{O}_{G/P}} \mathcal{O}_Z(-2u) - \text{mod} \rightarrow (\mathcal{D}_{G/P}^{(u+1)\omega_1+\omega_2} - \text{mod})^Z,$$

where $(\mathcal{D}_{G/P}^{(u+1)\omega_1+\omega_2} - \text{mod})^Z$ denotes the full subcategory of $\mathcal{D}_{G/P}^{(u+1)\omega_1+\omega_2}$ -mod supported on Z , and $\mathcal{O}_Z(2u)$ is the line bundle on Z with Chern class $2u$. Therefore we can put

$$J_{u,0} := \Gamma(\mathbb{P}^2, i_\star^u \mathcal{O}_Z(2u)).$$

It is clear that $J_{u,0}$ is a $(\mathfrak{g}, \mathfrak{k})$ -module, and furthermore, using the facts that $\mathcal{N} \simeq \mathcal{O}_Z(4)$ and that $i_\star^u \mathcal{O}_Z(2u)$ has a filtration with successive quotients $\mathcal{O}_Z(2u + 4(i+1))$, one easily verifies that $c(J_{u,0})$ is given by the right-hand side of (12). Since there are no finite-dimensional modules with central character $\chi(u+1, 1)$ for $u \in \frac{1}{2} + \mathbb{Z}$, $J_{u,0}$ is a simple \mathfrak{g} -module.

It remains to prove that every simple odd (D^u, \mathfrak{k}) -module is isomorphic to $J_{u,0}$ for some $u \in \frac{1}{2} + \mathbb{Z}$. Let M be a simple odd (D^u, \mathfrak{k}) -module and \hat{M} be a simple (A, \mathfrak{k}) -module such that $\text{Res}_u(\hat{M}) = M$. Then

by the proof of Lemma 5.14 $\hat{M}_f = 0$. For every $\mathfrak{b}_{\mathfrak{k}}$ -highest vector $v \in \text{Res}_u(\hat{M})$ there exists k such that $f^k \cdot v = 0$. Let v have weight m . Then by the relation $(d + \frac{5}{2})^2 = (\frac{m+1}{2})^2$ from the proof of Lemma 5.14, $\frac{m+1}{2} = \pm(u + 2k + \frac{5}{2})$, as $\mathbf{E}f^k \cdot v = (2k + u)f^k \cdot v$. Without loss of generality we may assume that m is very large and then $\frac{m+1}{2} = (u + 2k + \frac{5}{2})$. Therefore $\text{Hom}_{\mathfrak{k}}(V_m, M) \neq 0$ implies $m = 2u + 4k + 4$. Hence if M and M' are two odd (D^u, \mathfrak{k}) -modules one can find m such that $\text{Hom}_{\mathfrak{k}}(V_m, M) \neq 0$, $\text{Hom}_{\mathfrak{k}}(V_m, M') \neq 0$. But then $M \simeq M'$ by Corollary 5.8. \square

Let M be some A -module with semisimple \mathbf{E} -action. Consider the $U(\mathfrak{g})$ -modules $M^{(n)} := M \otimes S^n(\text{span}\{x, y, t\})$ for $n \in \mathbb{Z}_{\geq 0}$, together with the linear operators

$$\begin{aligned} \bar{d} : M^{(n)} &\rightarrow M^{(n-1)} \\ \bar{d} &= t \otimes \partial_t + x \otimes \partial_x + y \otimes \partial_y \\ \bar{\delta} : M^{(n)} &\rightarrow M^{(n+1)} \\ \bar{\delta} &= \partial_t \otimes t + \partial_x \otimes x + \partial_y \otimes y. \end{aligned}$$

It is straightforward to check that \bar{d} , $\mathbf{E} \otimes 1 - 1 \otimes \mathbf{E}$ and $\bar{\delta}$ form a standard $\mathfrak{sl}(2)$ -triple. Let $\text{Res}_s(M^{(k)})$ be the eigenspace of the operator $\mathbf{E} \otimes 1 + 1 \otimes \mathbf{E}$ in $M^{(k)}$. Then obviously \bar{d} and $\bar{\delta}$ induce operators

$$\begin{aligned} d : \text{Res}_s(M^{(n)}) &\rightarrow \text{Res}_s(M^{(n-1)}) \\ \delta : \text{Res}_s(M^{(n-1)}) &\rightarrow \text{Res}_s(M^{(n)}), \end{aligned}$$

and elementary $\mathfrak{sl}(2)$ representation theory implies that if $s \notin \mathbb{Z}$, $s < n - 1$ or $s \geq 2n$, then d is surjective, δ is injective, and

$$(13) \quad \text{Res}_s(M^{(n)}) = \ker d \oplus \text{im} \delta.$$

For any (D^u, \mathfrak{k}) -module M choose a simple $(A, \tilde{\mathfrak{k}})$ -module \hat{M} such that $\text{Res}_u(\hat{M}) = M$ (in fact \hat{M} is unique).

Let $T^n(M) := \text{Res}_{u+n}(\hat{M}^{(n)}) \cap \ker d$. If $u \neq -1, 0, \dots, n - 1$, (13) implies

$$(14) \quad c(T^n(M)) = c(\text{Res}_{u+n}(\hat{M}^{(n)})) - c(\text{Res}_{u+n}(\hat{M}^{(n-1)})).$$

Lemma 5.15. *Let M be a bounded simple (D^u, \mathfrak{k}) -module. Assume that $u \neq -1, 0, \dots, n - 1$. Then $T^n(M)$ is a simple $(\mathfrak{g}, \mathfrak{k})$ -module with central character $\chi(u + 1 - n, n + 1)$.*

Proof. Lemma 5.9 implies that M is a $(\mathfrak{g}, \mathfrak{k})$ -module with central character $\chi(u + 1, 1)$. Therefore $M \otimes S^n(\text{span}\{x, y, t\})$ has constituents with central character $\chi(u + 1 + n - 2k, 1 + k)$, $k = 0, \dots, n$, and $\text{im} \delta$ has constituents with central character $\chi(u + 1 + n - 2k, 1 + k)$, $k = 0, \dots, n - 1$.

Thus, $T^n(M)$ is a direct summand of $M \otimes S^n(\text{span}\{x, y, t\})$ with central character $\chi(u+1-n, n+1)$.

Our restrictions on u imply that the weights $(u+1)\omega_1 + \omega_2$ and $(u-n+1)\omega_1 + (n+1)\omega_2$ belong to the same Weyl chamber and have the same stabilizer in the Weyl group. Hence, T^n is nothing but the translation functor

$$T_{(u+1)\omega_1+\omega_2}^{(u-n+1)\omega_1+(n+1)\omega_2} : \mathfrak{B}_{\mathfrak{k}}^{\chi(u+1,1)} \rightarrow \mathfrak{B}_{\mathfrak{k}}^{\chi(u-n+1,n+1)}.$$

Therefore T^n is an equivalence of categories, in particular $T^n(M)$ is simple. \square

We put for $u \neq -1, 0, \dots, n-1$

$$I_{u,n}^{\pm} := T^n(I_{u,0}^{\pm}),$$

$$J_{u,n} := T^n(J_{u,0}).$$

Theorem 5.16. *Let M be a simple bounded infinite-dimensional $(\mathfrak{g}, \mathfrak{k})$ -module with central character χ . Then*

(a) if $\chi = \chi(u+1-n, n+1)$ for $u \notin \mathbb{Z}$,

$$M \simeq \begin{cases} I_{u,n}^{\pm} & \text{for } u \notin \frac{1}{2} + \mathbb{Z} \\ I_{u,n}^{\pm}, J_{u,n} & \text{for } u \in \frac{1}{2} + \mathbb{Z} \end{cases};$$

(b) if $\chi = \chi(u+1-n, n+1)$ for $u \in \mathbb{Z}_{\geq n}$,

$$M \simeq I_{-n-3, u-n}^{\pm}, I_{u,n}^{\pm};$$

(c) if $\chi = \chi(-1-n, n+1)$,

$$M \simeq I_{-2,n}^{\pm};$$

(d) if $\chi = \chi(0, n+1)$,

$$M \simeq (I_{-2,n}^{\pm})^{\tau},$$

where τ stands for the outer automorphism $\tau(X) = -X^t$ for any $X \in \mathfrak{g}$.

Proof. By Corollary 5.5 every simple bounded $(\mathfrak{g}, \mathfrak{k})$ -module has central character χ of the form $\chi(u+1-n, n+1)$ for some $n \in \mathbb{Z}_{\geq 0}$ and some $u \in \{\mathbb{C} \setminus \mathbb{Z}_{< n-1}\} \cup \{-2\}$. Moreover, $T^n = T_{(u+1)\omega_1+\omega_2}^{(u-n+1)\omega_1+(n+1)\omega_2}$ is an equivalence of the categories $\mathfrak{B}_{\mathfrak{k}}^{\chi(u+1,1)}$ and $\mathfrak{B}_{\mathfrak{k}}^{\chi(u+1-n, n+1)}$. If $u \notin \mathbb{Z}$, $\frac{1}{2} + \mathbb{Z}$ then $\mathfrak{B}_{\mathfrak{k}}^{\chi(u+1,1)}$ has two non-isomorphic simple objects, and, if $u \in \frac{1}{2} + \mathbb{Z}$, $\mathfrak{B}_{\mathfrak{k}}^{\chi(u+1,1)}$ has three non-isomorphic simple objects. This implies (a).

If $u \in \mathbb{Z}_{\geq 0}$, $u \geq n$, we have $\chi = \chi(u+1-n, n+1) = \chi((-n-3) + 1 - (u-n), (u-n) + 1)$, hence in this case $\mathfrak{B}_{\mathfrak{k}}^{\chi}$ has 4 non-isomorphic simple objects: $I_{u,n}^{\pm}$ and $I_{-n-3, u-n}^{\pm}$. This proves (b). If $n = -2$, $\mathfrak{B}_{\mathfrak{k}}^{\chi}$ is equivalent to $\mathfrak{B}_{\mathfrak{k}}^{\chi(1,1)}$ and has two simple objects, $I_{-2,n}^{\pm}$, which proves

(c). Finally if $u = n - 1$, the automorphism τ establishes an equivalence between $\mathfrak{B}_{\mathfrak{k}}^{\chi(0, n+1)}$ and $\mathfrak{B}_{\mathfrak{k}}^{\chi(-1-n, n+1)}$, hence (d). \square

Lemma 5.17. *For $a \in \mathbb{Z}_{\geq 2}$, define*

$$\mu_n(a, z) := \frac{z^a}{1 - z^4} \otimes c(V_{n,0}) - \frac{z^{a-2}}{1 - z^4} \otimes c(V_{n-1,0}).$$

For $a \in \mathbb{Z}_{\geq 0}$, define

$$\kappa_n(a, z) := \frac{z^a}{1 - z^4} \otimes c(V_{n,0}) - \frac{z^{a+2}}{1 - z^4} \otimes c(V_{n-1,0}).$$

Then

$$(15) \quad \mu_{2p}(a, z) = \frac{z^a}{1 - z^4} + \frac{z^{a-2}(z^4 + z^8 + \dots + z^{4p})}{1 - z^2},$$

$$(16) \quad \mu_{2p+1}(a, z) = \frac{z^a(1 + z^4 + \dots + z^{4p})}{1 - z^2},$$

$$(17) \quad \kappa_{2p}(a, z) = \frac{z^a}{1 - z^4} + \frac{z^{|a-4|} + \dots + z^{|a-4p|}}{1 - z^2},$$

$$(18) \quad \kappa_{2p+1}(a, z) = \frac{z^{|a-2|} + \dots + z^{|a-4p-2|}}{1 - z^2}.$$

Proof. Since $V_{n,0} = S^n(V_{1,0})$, and since $S^n(V_{1,0})$ is isomorphic as a \mathfrak{k} -module to $S^n(V_2)$, we have

$$c(V_{2p,0}) = 1 + z^4 + \dots + z^{2p},$$

$$c(V_{2p+1,0}) = z^2 + z^6 + \dots + z^{2p+2}.$$

Recall that $z^a \otimes z^b = \pi(z^a \sum_{i=0}^{i=b} z^{b-2i})$ (Lemma 3.2,(b)). Therefore

$$\begin{aligned} \frac{z^a}{1 - z^4} \otimes z^{2k} - \frac{z^{a-2}}{1 - z^4} \otimes z^{2k-2} &= \pi \left(\frac{z^{a-2} \left(z^2 \sum_{i=0}^{i=2k} z^{2k-2i} - \sum_{i=0}^{i=2k-2} z^{2k-2i} \right)}{1 - z^4} \right) = \\ &= \pi \left(\frac{z^{a-2} (z^{2k+2} + z^{2k})}{1 - z^4} \right) = \frac{z^{a-2+2k}}{1 - z^2}. \end{aligned}$$

$$\begin{aligned} \frac{z^a}{1-z^4} \otimes z^{2k} - \frac{z^{a+2}}{1-z^4} \otimes z^{2k-2} &= \pi \left(\frac{z^a \left(\sum_{i=0}^{2k} z^{2k-2i} - z^2 \sum_{i=0}^{2k-2} z^{2k-2-2i} \right)}{1-z^4} \right) = \\ &= \pi \left(\frac{z^a (z^{-2k} + z^{2-2k})}{1-z^4} \right) = \pi \left(\frac{z^{a-2k}}{1-z^2} \right) = \frac{z^{|a-2k|}}{1-z^2}. \end{aligned}$$

The above identities imply (15)-(18). \square

Theorem 5.18.

(a) Let $u \notin \mathbb{Z}, \frac{1}{2} + \mathbb{Z}$. Then

$$c(I_{u,n}^+) = \kappa_n(0, z), \quad c(I_{u,n}^-) = \mu_n(2, z).$$

(b) Let $u \in \frac{1}{2} + \mathbb{Z}$. Then

$$\begin{aligned} c(J_{u,n}) &= \kappa_n(4 + 2u, z) \quad \text{for } u \geq -\frac{1}{2}, \\ c(J_{u,n}) &= \mu_n(2 - 2u, z) \quad \text{for } u \leq -\frac{3}{2}. \end{aligned}$$

(c) Let $u \in 2\mathbb{Z}_{\geq 0}$. Then

$$\begin{aligned} c(I_{u,0}^+) &= \frac{z^{2u+4}}{1-z^4}, & c(I_{u,0}^-) &= \frac{z^2}{1-z^4}, \\ c(I_{u,n}^+) &= \kappa_n(2u + 4, z), & c(I_{u,n}^-) &= \mu_n(2, z). \end{aligned}$$

(d) Let $u \in 1 + 2\mathbb{Z}_{\geq 0}$. Then

$$\begin{aligned} c(I_{u,0}^+) &= \frac{1}{1-z^4}, & c(I_{u,0}^-) &= \frac{z^{2u+4}}{1-z^4}, \\ c(I_{u,n}^+) &= \kappa_n(0, z), & c(I_{u,n}^-) &= \kappa_n(2u + 4, z). \end{aligned}$$

(e) Let $u \in 2\mathbb{Z}_{\leq -2}$. Then

$$\begin{aligned} c(I_{u,0}^+) &= \frac{1}{1-z^4}, & c(I_{u,0}^-) &= \frac{z^{-2-2u}}{1-z^4}, \\ c(I_{u,n}^+) &= \kappa_n(0, z), & c(I_{u,n}^-) &= \mu_n(-2 - 2u, z). \end{aligned}$$

(f) Let $u \in -1 + 2\mathbb{Z}_{\leq -1}$. Then

$$\begin{aligned} c(I_{c,0}^+) &= \frac{z^{-2-2u}}{1-z^4}, & c(I_{u,0}^-) &= \frac{z^2}{1-z^4}, \\ c(I_{u,n}^+) &= \mu_n(-2 - 2u, z), & c(I_{u,n}^-) &= \mu_n(2, z). \end{aligned}$$

(g)

$$\begin{aligned} c(I_{-2,n}^+) &= c((I_{-2,n}^+)^{\tau}) = \kappa_n(0, z), \\ c(I_{-2,n}^-) &= c((I_{-2,n}^-)^{\tau}) = \mu_n(2, z). \end{aligned}$$

Proof. Using (14) one obtains the identities

$$(19) \quad \begin{aligned} c(I_{u,n}^{\pm}) &= c(I_{u,0}^{\pm} \otimes V_{n,0}) - c(I_{u+1,0}^{\mp} \otimes V_{n-1,0}), \\ c(J_{u,n}) &= c(J_{u,0} \otimes V_{n,0}) - c(J_{u+1,0} \otimes V_{n-1,0}). \end{aligned}$$

The theorem is a straightforward corollary of (19). Indeed, let us prove (f). In this case

$$\begin{aligned} c(I_{u,0}^+) &= \frac{z^{-2u-2}}{1-z^4}, & c(I_{u-1,0}^+) &= \frac{z^{-2u-4}}{1-z^4}, \\ c(I_{u,n}^+) &= \frac{z^{-2u-2}}{1-z^4} \otimes c(V_{n,0}) - \frac{z^{-2u-4}}{1-z^4} \otimes c(V_{n-1,0}) = \mu_n(-2-2u, z); \\ c(I_{u-1,0}^-) &= \frac{z^{-2u-4}}{1-z^4}, & c(I_{u-1,0}^+) &= \frac{1}{1-z^4}, \\ c(I_{u,n}^-) &= \frac{z^2}{1-z^4} \otimes c(V_{n,0}) - \frac{1}{1-z^4} \otimes c(V_{n-1,0}) = \mu_n(2, z). \end{aligned}$$

In all other cases the arguments are similar. \square

Corollary 5.19.

- (a) *The minimal \mathfrak{k} -type can be any V_k but its multiplicity is always 1.*
- (b) *For sufficiently large i $c_i(M) = c_{i+4}(M)$ for any simple bounded $(\mathfrak{g}, \mathfrak{k})$ -module, and for sufficiently large j there are the following \mathfrak{k} -multiplicities:*

$$\begin{aligned} c_{4j}(I_{u,2p+1}^\pm) &= c_{4j+2}(I_{u,2p+1}^\pm) = p+1, \\ c_{4j}(I_{u,2p}^+) &= p+1, c_{4j+2}(I_{u,2p}^+) = p, \\ c_{4j+2}(I_{u,2p}^-) &= p+1, c_{4j}(I_{u,2p}^-) = p, \\ c_{4j+1}(J_{u,2p+1}) &= c_{4j+3}(J_{u,2p+1}) = p+1, \\ c_{4j+2u}(J_{u,2p}) &= p, c_{4j+2u+2}(J_{u,2p}) = p+1. \end{aligned}$$

- (c) *The only multiplicity-free simple infinite-dimensional $(\mathfrak{g}, \mathfrak{k})$ -modules are $I_{u,0}^\pm, J_{u,0}, I_{u,1}^\pm, J_{u,1}, (I_{-2,1}^\pm)^\tau$.*

The complete list of multiplicity-free simple $(\mathfrak{g}, \mathfrak{k})$ -modules has been first found by Dj. Sijacki, see [S] and the references therein for a historic perspective on this problem.

6. CLASSIFICATION OF SIMPLE BOUNDED $(\mathfrak{sp}(4), \mathfrak{sl}(2))$ -MODULES

In this section we classify all simple bounded $(\mathfrak{g}, \mathfrak{k})$ -modules, where $\mathfrak{g} = \mathfrak{sp}(4)$ and \mathfrak{k} is a principal $\mathfrak{sl}(2)$ -subalgebra or a $\mathfrak{sl}(2)$ -subalgebra corresponding to a short root. We fix a Cartan subalgebra $\mathfrak{h} \subset \mathfrak{g}$ and write the roots of \mathfrak{g} as $\{\pm 2\epsilon_1, \pm 2\epsilon_2, \pm\epsilon_1 \pm \epsilon_2\}$. Our fixed simple roots are $\epsilon_1 - \epsilon_2, 2\epsilon_2$, and $\rho = 2\epsilon_1 + \epsilon_2$. By $e_1, e_2, h_1, h_2, f_1, f_2$ we denote the Serre generators of \mathfrak{g} associated to our choice of simple roots, [OV]. We define two $\mathfrak{sl}(2)$ -subalgebras of \mathfrak{g} : one with basis e_1, h_1, f_1 and one with basis $e_1 + 2e_2, 3h_1 + 4h_2, 3f_1 + 2f_2$. The first one is the root subalgebra corresponding to the simple root $\epsilon_1 - \epsilon_2$, and the second one is a principal $\mathfrak{sl}(2)$ -subalgebra. In Sections 6 and 7, we denote by

\mathfrak{k} any one of these two subalgebras, referring respectively to the *root case* and to the *principal case* when we want to be specific. We set $\mathfrak{b}_{\mathfrak{k}} := \mathfrak{b} \cap \mathfrak{k}$, where \mathfrak{b} is the Borel subalgebra generated by e_1, e_2, h_1, h_2 . By $L_{a,b}$ we denote the simple \mathfrak{b} -highest weight \mathfrak{g} -module with highest weight $a\epsilon_1 + b\epsilon_2 - \rho = (a-2)\epsilon_1 + (b-1)\epsilon_2$, by $V_{a,b}$ we denote the simple finite-dimensional \mathfrak{g} -module with highest weight $a\epsilon_1 + b\epsilon_2$, and $\chi(a, b)$ is the central character of $L_{a,b}$.

Lemma 6.1. *Let $\dim L_{a,b} = \infty$ and $\text{GKdim} L_{a,b} \leq 2$. Then $a > |b|$ and $a, b \in \frac{1}{2} + \mathbb{Z}$.*

Proof. Let $\lambda = a\epsilon_1 + b\epsilon_2$. If $(\lambda, \alpha) \notin \mathbb{Z}_{>0}$ for all positive roots α , then $L_{a,b}$ is a Verma module and therefore its Gelfand-Kirillov dimension equals 4. If $(\lambda, \check{\alpha}) \in \mathbb{Z}_{>0}$ for exactly one positive root, then one has the following exact sequence

$$0 \rightarrow L_{w_{\alpha}(\lambda)} \rightarrow M_{\lambda} \rightarrow L_{\lambda} \rightarrow 0,$$

where w_{α} denotes the reflection in α . A straightforward computation shows that in this case $\text{GKdim} L_{\lambda} = 3$. Therefore $\text{GKdim} L_{\lambda} \leq 2$ implies the existence of two positive roots α and β such that $(\lambda, \check{\alpha}), (\lambda, \check{\beta}) \in \mathbb{Z}_{>0}$. One can see immediately that at least one of these roots, say α , is simple. If N_{λ} denotes the quotient of M_{λ} by the submodule generated by a highest vector with weight $w_{\alpha}(\lambda) - \rho$, then $\text{GKdim} N_{\lambda} = 3$. The condition $\text{GKdim} L_{\lambda} \leq 2$ implies the reducibility of N_{λ} which in turn implies $(\lambda, \check{\gamma}) \in \mathbb{Z}_{>0}$ for the positive root γ orthogonal to α . That leaves only two possibilities for λ : λ is either regular integral or λ satisfies the conditions of the Lemma.

It remains to eliminate the case of a regular integral non-dominant λ . By using the translation functor we may assume without loss of generality that λ belongs to the Weyl group orbit of ρ . That leaves four possibilities for λ : $2\epsilon_1 - \epsilon_2, \epsilon_1 - 2\epsilon_2, \epsilon_1 + 2\epsilon_2, -\epsilon_1 + 2\epsilon_2$. Let \mathfrak{p}_1 and \mathfrak{p}_2 be the parabolic subalgebras obtained from \mathfrak{b} by joining $\epsilon_2 - \epsilon_1$ and $-2\epsilon_2$ respectively. It is not difficult to verify the existence of embeddings

$$\begin{aligned} L_{2,-1} &\rightarrow U(\mathfrak{g}) \otimes_{U(\mathfrak{p}_1)} F_{2,1}^1, & L_{1,-2} &\rightarrow U(\mathfrak{g}) \otimes_{U(\mathfrak{p}_1)} F_{2,-1}^1, \\ L_{1,2} &\rightarrow U(\mathfrak{g}) \otimes_{U(\mathfrak{p}_2)} F_{2,1}^2, & L_{-1,2} &\rightarrow U(\mathfrak{g}) \otimes_{U(\mathfrak{p}_2)} F_{1,2}^2, \end{aligned}$$

where $F_{a,b}^1$ (respectively, $F_{a,b}^2$) is the finite dimensional \mathfrak{p}_1 -module (resp., \mathfrak{p}_2 -module) with \mathfrak{b} -highest weight $a\epsilon_1 + b\epsilon_2 - \rho$. Therefore the Gelfand-Kirillov dimension of any of the above four simple modules equals the Gelfand-Kirillov dimension of the corresponding parabolically induced module, i.e. 3. The proof is now complete. \square

Corollary 6.2. *Let M be a simple bounded infinite-dimensional $(\mathfrak{g}, \mathfrak{k})$ -module. Then $\text{Ann}M = \text{Ann}L_{a,b}$ for some a, b with $a > |b|$, $a, b \in \frac{1}{2} + \mathbb{Z}$. In particular, $\chi(a, b)$ is the central character of M .*

Proof. By Duflo's theorem, $\text{Ann}M = \text{Ann}L_{a,b}$ for some a, b . It is known that $\frac{1}{2} \dim X_{L_{a,b}} = \text{GKdim}L_{a,b}$, thus $\text{GKdim}M \geq \text{GKdim}L_{a,b}$. On the other hand, $\text{GKdim}M \leq 2 = b_{\mathfrak{k}}$ holds by Theorem 4.4 in [PS2]. Hence $\text{GKdim}L_{a,b} \leq 2$, and Lemma 6.1 applies to $L_{a,b}$. \square

Corollary 6.3. *Let $a, b \in \frac{1}{2} + \mathbb{Z}$, $a > |b|$. Then $\mathfrak{B}_{\mathfrak{k}}^{\chi(a,b)}$ is equivalent to $\mathfrak{B}_{\mathfrak{k}}^{\chi(\frac{3}{2}, \frac{1}{2})}$.*

Proof. It is well known that the categories $U^{\chi(a,b)}\text{-mod}$ for (a, b) as above are translation-equivalent to the category $U^{\chi(\frac{3}{2}, \frac{1}{2})}\text{-mod}$. Since the translation functor preserves the subcategories of bounded modules, the categories $\mathfrak{B}_{\mathfrak{k}}^{\chi(a,b)}$ and $\mathfrak{B}_{\mathfrak{k}}^{\chi(\frac{3}{2}, \frac{1}{2})}$ are equivalent as well. \square

Our next step is to describe the quotient algebra $U(\mathfrak{g})/\text{Ann}L_{\frac{3}{2}, \frac{1}{2}}$. In this section we denote by A the Weyl algebra in two variables, i.e. the algebra of differential operators acting in $\mathbb{C}[x, y]$. We introduce a \mathbb{Z}_2 -grading, $A := A_0 \oplus A_1$, by putting $\deg x = \deg y = \deg \partial_x = \deg \partial_y := \bar{1} \in \mathbb{Z}_2$. It is well known that there exists a surjective algebra homomorphism

$$\kappa : U(\mathfrak{g}) \rightarrow A_0$$

such that

$$\kappa(e_1) = x\partial_y, \quad \kappa(e_2) = \frac{y^2}{2}, \quad \kappa(f_1) = y\partial_x, \quad \kappa(f_2) = -\frac{\partial_y^2}{2},$$

$$\kappa(h_1) = x\partial_x - y\partial_y, \quad \kappa(h_2) = y\partial_y + \frac{1}{2}.$$

The kernel of κ equals $\text{Ann}L_{\frac{3}{2}, \frac{1}{2}}$. Furthermore, $\kappa(\mathfrak{k})$ is spanned by $E := x\partial_y$, $F := y\partial_x$, $H := x\partial_x - y\partial_y$ in the root case, and respectively by $E := x\partial_y + y^2$, $H := 3x\partial_x + y\partial_y + 2$, $F := 3y\partial_x - \partial_y^2$ in the principal case.

The problem of describing all simple modules in $\mathfrak{B}_{\mathfrak{k}}^{\chi(\frac{3}{2}, \frac{1}{2})}$ is equivalent to the problem of describing all simple (A_0, \mathfrak{k}) -modules, i.e. all simple locally $\kappa(\mathfrak{k})$ -finite A_0 -modules. The following lemma reduces this problem to a classification of all simple (A, \mathfrak{k}) -modules.

Lemma 6.4. *Every simple (A, \mathfrak{k}) -module M is a \mathbb{Z}_2 -graded A -module, i. e. $M = M_0 \oplus M_1$ where M_0 and M_1 are simple (A_0, \mathfrak{k}) -modules. Furthermore, $M = A \otimes_{A_0} M_0$, and the \mathbb{Z}_2 -grading on M is unique up to interchanging M_0 with M_1 .*

Proof. The element H (as defined above separately for the root case and for the principal case) acts semisimply on M with integer eigenvalues. We define M_0 (respectively, M_1) as the direct sum of H -eigenspaces with even (resp., odd) eigenvalues. It is obvious that $M = M_0 \oplus M_1$, that M_0 and M_1 are simple A_0 modules, and that $M = A \otimes_{A_0} M_0$. Since M_0 and M_1 are non-isomorphic as A_0 -modules, the uniqueness follows from the fact that a decomposition of M as an A_0 -module into a direct sum of two non-isomorphic A_0 -modules is unique. \square

Remark. More generally, if \mathfrak{k}' is a subalgebra of $\mathfrak{g}' = \mathfrak{sp}(2m)$ such that the centralizer of \mathfrak{k}' in the Weyl A' algebra of m indeterminates is abelian, every (A', \mathfrak{k}') -module is a multiplicity-free $(\mathfrak{g}', \mathfrak{k}')$ -module whose primitive ideal is a Joseph ideal. F. Knop has classified all such subalgebras \mathfrak{k}' , [Kn2], which makes us optimistic that this idea can eventually lead to a classification of simple bounded $(\mathfrak{g}', \mathfrak{k}')$ -modules.

Let $\text{Fou} : A \rightarrow A$ be the automorphism defined by

$$\text{Fou}(x) := \partial_x, \quad \text{Fou}(y) := \partial_y, \quad \text{Fou}(\partial_x) := -x, \quad \text{Fou}(\partial_y) := -y$$

If M is an A -module, we denote by M^{Fou} the twist of M by Fou .

Theorem 6.5. *In the root case, any simple (A, \mathfrak{k}) -module is isomorphic to $\mathbb{C}[x, y]$ or $\mathbb{C}[x, y]^{\text{Fou}}$.*

Proof. Let M be a simple (A, \mathfrak{k}) -module. Then there exists $0 \neq v \in M$ such that $E \cdot v = 0$, i.e. $x\partial_y \cdot v = 0$. Hence either x or ∂_y act locally nilpotently on M .

Assume first that ∂_y acts locally nilpotently on M . Then $\partial_x \in [\mathfrak{k}, \partial_y]$ also acts locally nilpotently on M . Let A^+ be the abelian subalgebra in A generated by ∂_x, ∂_y . One can find $0 \neq w \in M$ such that $A^+ \cdot w = 0$, and hence

$$M \cong A \otimes_{A^+} \mathbb{C} \cong \mathbb{C}[x, y].$$

If x acts locally nilpotently on M , one considers M^{Fou} and reduces to the previous case. \square

Corollary 6.6. *In the root case, up to isomorphism, there are exactly four simple $(\mathfrak{g}, \mathfrak{k})$ -modules with central character $\chi(\frac{3}{2}, \frac{1}{2})$. As \mathfrak{k} -modules two of these modules are isomorphic to*

$$V_0 \oplus V_2 \oplus V_4 \oplus \dots \quad ,$$

and the other two are isomorphic to

$$V_1 \oplus V_3 \oplus V_5 \oplus \dots \quad .$$

Theorem 6.7. *In the principal case, up to isomorphism, there exist exactly two simple (A, \mathfrak{k}) -modules and they have the following \mathfrak{k} -module decompositions:*

$$V_0 \oplus V_3 \oplus V_6 \oplus V_9 \oplus \dots, \quad V_1 \oplus V_4 \oplus V_7 \oplus V_{10} \oplus \dots$$

Proof. Note that \mathfrak{k} is a maximal subalgebra of \mathfrak{g} . Hence, every element $g \in \mathfrak{g} \setminus \mathfrak{k}$ acts freely on a simple (A, \mathfrak{k}) -module M . In particular, x^2 acts freely on M , and therefore x acts freely on M . Let A_x be the localization of A in x , and $M_x := A_x \otimes_A M$. Then $M \subset M_x$. Fix $0 \neq m \in M$ with $E \cdot m = 0$ and $H \cdot m = \lambda m$ for a minimal $\lambda \in \mathbb{Z}_{\geq 0}$. Since $E = x\partial_y + y^2$ and $H = 3x\partial_x + y\partial_y + 2$, we have

$$\partial_y \cdot m = -\frac{y^2}{x} \cdot m, \quad \partial_x \cdot m = \left(\frac{y^3}{3x^2} + \frac{\lambda - 2}{3x} \right) \cdot m.$$

Therefore, $M_x = \mathbb{C}[x, x^{-1}, y] \cdot m$. Set

$$u_\lambda := x^{\frac{\lambda-2}{3}} \exp\left(\frac{-y^3}{3x}\right).$$

Then it is easy to see that M_x is isomorphic to $\mathcal{F}_\lambda := \mathbb{C}[x, x^{-1}, y]u_\lambda$ and that $\mathcal{F}_\lambda = \mathcal{F}_{\lambda+3}$. Hence, M_x is isomorphic to $\mathcal{F}_0, \mathcal{F}_1$ or \mathcal{F}_2 .

Next we calculate $\Gamma_{\mathfrak{k}}(\mathcal{F}_\lambda)$. Note that the space of $\mathfrak{b}_{\mathfrak{k}}$ -singular vectors in \mathcal{F}_λ is spanned by the family $u_{\lambda+3k}$, $k \in \mathbb{Z}$ of solutions to the differential equation

$$E \cdot u = x\partial_y(u) + y^2u = 0.$$

If $\lambda \in \mathbb{Z}_{\geq 0}$, then $F^{\lambda+1} \cdot u_\lambda$ is again a $\mathfrak{b}_{\mathfrak{k}}$ -highest vector of weight $-\lambda - 2$. Therefore $F^{\lambda+1} \cdot u_\lambda = cu_{-\lambda-2}$ for some constant c . On the other hand, $u_{-\lambda-2} \in \mathcal{F}_\lambda$ iff $\lambda - (-\lambda - 2) = 2\lambda + 2 \in 3\mathbb{Z}$ or $\lambda = 3k + 2$. Hence $F^{\lambda+1} \cdot u_\lambda = 0$ for $\lambda = 3k$ or $\lambda = 3k + 1$. Thus, $\Gamma_{\mathfrak{k}}(\mathcal{F}_0)$ is generated by u_{3k} for $k \geq 0$, $\Gamma_{\mathfrak{k}}(\mathcal{F}_1)$ is generated by u_{3k+1} for $k \geq 0$, and we have the \mathfrak{k} -module decompositions

$$\Gamma_{\mathfrak{k}}(\mathcal{F}_0) \simeq V_0 \oplus V_3 \oplus V_6 \oplus V_9 \oplus \dots, \quad \Gamma_{\mathfrak{k}}(\mathcal{F}_1) \simeq V_1 \oplus V_4 \oplus V_7 \oplus V_{10} \oplus \dots$$

Let us prove that $\Gamma_{\mathfrak{k}}(\mathcal{F}_0)$ and $\Gamma_{\mathfrak{k}}(\mathcal{F}_1)$ are simple A -modules. Indeed, let N be a proper submodule of $\Gamma_{\mathfrak{k}}(\mathcal{F}_0)$. If $u_\lambda \in N$, then $u_{\lambda+3k} = x^k u_\lambda \in N$ for all positive k . Choose the minimal λ such that $u_\lambda \in N$. Then the quotient module has a decomposition $V_{\lambda-3} \oplus \dots \oplus V_0$, hence it is finite-dimensional. Since A has no non-zero finite-dimensional modules, this is a contradiction. The case of $\Gamma_{\mathfrak{k}}(\mathcal{F}_1)$ is very similar. In this way we obtain that, if $M_x = \mathcal{F}_0$ or \mathcal{F}_1 , then M is respectively isomorphic to $\Gamma_{\mathfrak{k}}(\mathcal{F}_0)$ or $\Gamma_{\mathfrak{k}}(\mathcal{F}_1)$.

Finally, we show that $\Gamma_{\mathfrak{k}}(\mathcal{F}_2) = 0$. It is sufficient to check that there is no non-zero $v \in \mathcal{F}_2$ with $F \cdot v = 0$ and

$$(20) \quad H \cdot v = (-3k - 2)v \text{ for } k \in \mathbb{Z}_{\geq 0}.$$

Indeed, then v would be a solution of the differential equation

$$3yv_x = v_{yy}.$$

Since $v \in \mathcal{F}_2$,

$$v = g(x, y) \exp\left(-\frac{y^3}{3x}\right)$$

for some $g(x, y) \in \mathbb{C}[x, x^{-1}, y]$ such that

$$3yg_x = g_{yy} - 2\frac{y^2}{x}g_y - 2\frac{y}{x}g.$$

As $g(x, y)$ is homogeneous with respect to H , we may assume without loss of generality that

$$g(x, y) = \sum_{i=0}^l b_i x^{p-i} y^{3i+s},$$

where $s \in \mathbb{Z}_{\geq 0}$, $p \in \mathbb{Z}$, $b_i \in \mathbb{C}$, $b_0 = 1$. The equation on the highest term with respect to x gives the condition

$$\partial_y^2 (y^s) = 0,$$

or, equivalently, $s = 0, 1$. But $H \cdot g = (3p + s + 2)g$, hence $H \cdot v = (3p + s + 2) \cdot v$. Therefore

$$H \cdot v = (3p + 2)v \text{ or } H \cdot v = (3p + 3)v,$$

and (20) does not hold. \square

Theorem 6.7 together with Lemma 6.4 yield the following.

Corollary 6.8. *In the principal case, up to isomorphism, there are exactly four simple $(\mathfrak{g}, \mathfrak{k})$ -modules with central character $\chi(\frac{3}{2}, \frac{1}{2})$. They have the following \mathfrak{k} -module decompositions:*

$$(21) \quad V_0 \oplus V_6 \oplus V_{12} \oplus \dots, V_1 \oplus V_7 \oplus V_{13} \oplus \dots, V_3 \oplus V_9 \oplus V_{15} \oplus \dots, V_4 \oplus V_{10} \oplus V_{16} \oplus \dots$$

7. \mathfrak{k} -CHARACTERS OF SIMPLE BOUNDED $(\mathfrak{sp}(4), \mathfrak{sl}(2))$ -MODULES

7.1. The root case. In this case, the four simple modules of Corollary 6.6 are nothing but the simple highest weight modules $L_{\frac{3}{2}, \frac{1}{2}}$, $L_{\frac{3}{2}, -\frac{1}{2}}$, and their respective restricted duals $L'_{\frac{3}{2}, \frac{1}{2}}$, $L'_{\frac{3}{2}, -\frac{1}{2}}$, i.e. the simple \mathfrak{b} -lowest weight modules with lowest weights $(-\frac{3}{2}, -\frac{1}{2})$ and $(-\frac{3}{2}, \frac{1}{2})$. Therefore,

by Corollaries 6.2, 6.3 we conclude that all simple bounded $(\mathfrak{g}, \mathfrak{k})$ -modules are precisely $L_{a,b}$ and the lowest weight modules $L'_{-a,-b}$, where $a > |b| \in \frac{1}{2} + \mathbb{Z}$. Since $c(L_{a,b}) = c(L'_{-a,-b})$, it suffices to compute $c(L_{a,b})$, for a, b as above.

The \mathfrak{h} -character of $L_{a,b}$ is given by the formula

$$(22) \quad \text{ch}_{\mathfrak{h}} L_{a,b} = \frac{(x^{a-b} - x^{b-a})(y^{a+b} - y^{-a-b})}{(x - x^{-1})(y - y^{-1})(xy - x^{-1}y^{-1})(x^{-1}y - xy^{-1})},$$

where $x = e^{\frac{\epsilon_1 - \epsilon_2}{2}}$, $y = e^{\frac{\epsilon_1 + \epsilon_2}{2}}$. We rewrite (22) as

$$(23) \quad \frac{(x^{a-b} - x^{b-a})(y^{a-b} - y^{b-a})}{(x - x^{-1})(y - y^{-1})} y^{-2} (1 - x^2 y^{-2})^{-1} (1 - x^{-2} y^{-2})^{-1}.$$

Next we note that

$$(24) \quad (1 - x^2 y^{-2})^{-1} (1 - x^{-2} y^{-2})^{-1} = \sum_{k=0}^{\infty} y^{-2k} (x^{2k} + x^{2k-4} + \dots + x^{-2k}),$$

and use the expression

$$z^k = x^k + x^{k-2} + \dots + x^{-k} = \frac{x^{k+1} - x^{-(k+1)}}{x - x^{-1}}$$

to rewrite the right-hand side of (24) in the form

$$\sum_{k=0}^{\infty} y^{-2k} (z^{2k} - z^{2k-2} + \dots + (-1)^k) = \frac{1}{1 + y^2} \sum_{k=0}^{\infty} z^{2k} y^{-2k}.$$

Now (23) becomes

$$\text{ch}_{\mathfrak{h}} L_{a,b} = z^{a-b-1} \frac{y^{a+b} - y^{-a-b}}{y - y^{-1}} \frac{1}{1 + y^2} \sum_{k=0}^{\infty} z^{2k} y^{-2k}.$$

To find the \mathfrak{k} -character of $L_{a,b}$, we set $y = 1$:

$$(25) \quad c(L_{a,b}) = \frac{a+b}{2} z^{a-b-1} \otimes \sum_{k=0}^{\infty} z^{2k}.$$

Thus, equation (25) implies the following result.

Theorem 7.1.

(a) *If $a - b$ is even and $a + b$ is odd, then*

$$c(L_{a,b}) = \frac{a+b}{2} (2z + 4z^3 + \dots + (a-b)z^{a-b-1} + (a-b)z^{a-b+1} + \dots).$$

(b) *If $a - b$ is odd and $a + b$ is even, then*

$$c(L_{a,b}) = \frac{a+b}{2} (1 + 3z^2 + 5z^4 + \dots + (a-b)z^{a-b-1} + (a-b)z^{a-b+1} + \dots).$$

- (c) In the case (a) the minimal \mathfrak{k} -type is V_1 and its multiplicity is $a+b$. In the case (b) the minimal \mathfrak{k} -type is V_0 and its multiplicity is $\frac{a+b}{2}$.
- (d) For sufficiently large i ,

$$c_i(L_{a,b}) = c_{i+2}(L_{a+b}) = \frac{(a^2 + b^2)(1 + (-1)^{a+b-i})}{4}.$$

- (e) $L_{a,b}$ is \mathfrak{k} -multiplicity-free if and only if $a = \frac{3}{2}$, hence the only simple multiplicity-free $(\mathfrak{g}, \mathfrak{k})$ -modules are those with central character $\chi(\frac{3}{2}, \frac{1}{2})$, i.e. the four \mathfrak{g} -modules from Corollary 6.8.

7.2. The principal case. We now proceed to calculating the \mathfrak{k} -characters of all simple bounded $(\mathfrak{g}, \mathfrak{k})$ -modules where $\mathfrak{g} = \mathfrak{sp}(4)$ and \mathfrak{k} is the principal subalgebra of \mathfrak{g} fixed in Section 6. In this case, let $M_{\frac{3}{2}, \frac{1}{2}}^0$ and $M_{\frac{3}{2}, \frac{1}{2}}^1$ denote the simple bounded $(\mathfrak{g}, \mathfrak{k})$ -modules with central character $\chi(\frac{3}{2}, \frac{1}{2})$ and respective \mathfrak{k} -module decompositions $V_0 \oplus V_6 \oplus V_{12} \oplus \dots$ and $V_1 \oplus V_7 \oplus V_{13} \oplus \dots$. We set $M_{a,b}^s := T_{a\epsilon_1 + b\epsilon_2}^{\frac{3}{2}\epsilon_1 + \frac{1}{2}\epsilon_2}(M_{\frac{3}{2}, \frac{1}{2}}^s)$ for $a, b \in \frac{1}{2} + \mathbb{Z}$, $a > |b|$, $s \in \{0, 1\}$, and $M_{a,b}^s := 0$ for $a, b \in \frac{1}{2} + \mathbb{Z}$, $a \leq |b|$, $s \in \{0, 1\}$. By $V_{p,q}$ we denote the simple finite-dimensional $\mathfrak{g} = \mathfrak{sp}(4)$ -module with \mathfrak{b} -highest weight $p\epsilon_1 + q\epsilon_2$ ($p, q \in \mathbb{Z}_{\geq 0}$, $p \geq q$).

Lemma 7.2. *We have*

$$(26) \quad V_{1,0} \otimes M_{a,b}^s \simeq M_{a+1,b}^s \oplus M_{a,b+1}^s \oplus M_{a-1,b}^s \oplus M_{a,b-1}^s,$$

and, for $a \neq |b| + 1$,

$$(27) \quad V_{1,1} \otimes M_{a,b}^s \simeq M_{a+1,b+1}^s \oplus M_{a,b}^s \oplus M_{a-1,b+1}^s \oplus M_{a+1,b-1}^s \oplus M_{a-1,b-1}^s.$$

If $a = b + 1$, $b > 0$, then

$$(28) \quad V_{1,1} \otimes M_{a,b}^s \simeq M_{a+1,b+1}^s \oplus M_{a+1,b-1}^s \oplus M_{a-1,b-1}^s,$$

and if $a = -b + 1$, $b < 0$, then

$$(29) \quad V_{1,1} \otimes M_{a,b}^s \simeq M_{a+1,b+1}^s \oplus M_{a+1,b-1}^s \oplus M_{a-1,b+1}^s.$$

Proof. Let us first prove (26). In what follows we use the notation of [PS2], Section 5. Let $\mathcal{M}_{a,b}^s := \mathcal{D}_{G/B}^{a,|b|} \otimes_{U\chi(a,b)} M_{a,b}^s$ be the localization of $M_{a,b}$ on G/B . Then as a sheaf of U -modules $V_{1,0} \otimes \mathcal{M}_{a,b}^s$ has a filtration of length 4 with the following associated factors given in increasing order:

$$\mathcal{O}(-\epsilon_1) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s, \quad \mathcal{O}(-\epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s, \quad \mathcal{O}(\epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s, \quad \mathcal{O}(\epsilon_1) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s.$$

Note that Z_U acts via a character on any of the four associated factors, and that these characters are pairwise distinct. Therefore, as a sheaf of U -modules, $V_{1,0} \otimes \mathcal{M}_{a,b}^s$ is isomorphic to the direct sum

$$(\mathcal{O}(-\epsilon_1) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s) \oplus (\mathcal{O}(-\epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s) \oplus (\mathcal{O}(\epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s) \oplus (\mathcal{O}(\epsilon_1) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s).$$

Now we calculate $\Gamma(G/B, V_{1,0} \otimes \mathcal{M}_{a,b}^s)$. If $a = b + 1, b > 0$, then

$$\Gamma(G/B, \mathcal{O}(-\epsilon_1) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s) = \Gamma(G/B, \mathcal{O}(\epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s) = 0$$

as there are no bounded modules with these central characters. Similarly, if $a = -b + 1, b < 0$, then

$$\Gamma(G/B, \mathcal{O}(-\epsilon_1) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s) = \Gamma(G/B, \mathcal{O}(-\epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s) = 0.$$

In all other cases

$$\Gamma(G/B, \mathcal{O}(\pm\epsilon_1) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s) \simeq M_{a\pm 1, b}^s,$$

$$\Gamma(G/B, \mathcal{O}(\pm\epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s) \simeq M_{a, b\pm 1}^s.$$

Thus, (26) is established.

Consider (27). Then as a sheaf of U -modules $V_{1,1} \otimes \mathcal{M}_{a,b}^s$ has a filtration of length 5 with the following associated factors given in increasing order:

$$\begin{aligned} &\mathcal{O}(-\epsilon_1 - \epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s, \quad \mathcal{O}(\epsilon_1 - \epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s, \quad \mathcal{M}_{a,b}^s, \\ &\mathcal{O}(-\epsilon_1 + \epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s, \quad \mathcal{O}(\epsilon_1 + \epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s. \end{aligned}$$

Note that Z_U acts via a character on any of the five associated factors, and that these characters are pairwise distinct if $a \neq |b| + 1$. Therefore the proof of (27) is very similar to that of (26).

Let now $a = b + 1$. Then $\mathcal{M}_{a,b}^s$ and $\mathcal{O}(-\epsilon_1 + \epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s$ both afford the central character $\chi(a, b)$. Thus, as a sheaf of U -modules, $V_{1,1} \otimes \mathcal{M}_{a,b}^s$ is isomorphic to the direct sum

$$(30) \quad (\mathcal{O}(-\epsilon_1 - \epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s) \oplus (\mathcal{O}(\epsilon_1 - \epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s) \oplus (\mathcal{M}_{a,b}^s)' \oplus \\ \oplus (\mathcal{O}(\epsilon_1 + \epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s),$$

where for $(\mathcal{M}_{a,b}^s)'$ we have an exact sequence

$$0 \rightarrow \mathcal{M}_{a,b}^s \rightarrow (\mathcal{M}_{a,b}^s)' \rightarrow \mathcal{O}(-\epsilon_1 + \epsilon_2) \otimes_{\mathcal{O}} \mathcal{M}_{a,b}^s \rightarrow 0.$$

We will show that $\Gamma(G/B, (\mathcal{M}_{a,b}^s)') = 0$. It suffices to show that the tensor product $V_{1,1} \otimes \mathcal{M}_{a,b}^s$ has no simple constituent with central character $\chi(a, b)$. Indeed, from (26), we see that $V_{1,0} \otimes V_{1,0} \otimes \mathcal{M}_{a,b}^s$ has exactly two simple constituents affording the central character $\chi(a, b)$ and that both these constituents are isomorphic to $M_{a,b}^s$. Recall that

$$V_{1,0} \otimes V_{1,0} \cong V_{2,0} \oplus V_{1,1} \oplus V_{0,0}.$$

Clearly, $V_{0,0} \otimes M_{a,b}^s = M_{a,b}^s$. Furthermore, $V_{2,0}$ is the adjoint representation and therefore the very \mathfrak{g} -module structure on $M_{a,b}^s$ defines a non-trivial intertwining operator $V_{2,0} \otimes M_{a,b}^s \rightarrow M_{a,b}^s$. Thus, $V_{2,0} \otimes M_{a,b}^s$ must have a constituent isomorphic to $M_{a,b}^s$ and consequently $V_{1,1} \otimes M_{a,b}^s$ has no simple constituent affording the central character $\chi(a,b)$. By taking the global sections of the direct sum (30) we obtain (28). The case $a = -b + 1$, which leads to (29), is similar. \square

Lemma 7.3. *There is the following \mathfrak{k} -module decomposition*

$$(31) \quad M_{\frac{3}{2}, -\frac{1}{2}}^s \simeq V_{3+s} \oplus V_{9+s} \oplus V_{15+s} \oplus \dots$$

Proof. By (26),

$$M_{\frac{3}{2}, \frac{1}{2}}^0 \otimes V_{1,0} \simeq M_{\frac{5}{2}, \frac{1}{2}}^0 \oplus M_{\frac{3}{2}, -\frac{1}{2}}^0.$$

As a \mathfrak{k} -module, $V_{1,0}$ is isomorphic to V_3 . Hence $M_{\frac{3}{2}, \frac{1}{2}}^0 \otimes V_{1,0}$ has a \mathfrak{k} -module decomposition

$$2V_3 \oplus V_5 \oplus \dots$$

Since $\chi(\frac{3}{2}, -\frac{1}{2}) = \chi(\frac{3}{2}, \frac{1}{2})$, $M_{\frac{3}{2}, -\frac{1}{2}}^0$ must have one of the four \mathfrak{k} -module decompositions (21), and hence (26) implies (31) for $s = 0$. Similarly, $M_{\frac{3}{2}, \frac{1}{2}}^1 \otimes V_{1,0}$ has the \mathfrak{k} -module decomposition $V_2 \oplus 2V_4 \oplus \dots$, which implies (31) for $s = 1$. \square

We set now $\varphi_{a,b}^s(z) := c(M_{a,b}^s)$ for $a, b \in \frac{1}{2} + \mathbb{Z}, a \geq |b|, s \in \{0, 1\}$ and extend the definition of $\varphi_{a,b}^s(z)$ to arbitrary pairs $a, b \in \frac{1}{2} + \mathbb{Z}$ by putting

$$(32) \quad \varphi_{a,b}^s(z) = -\varphi_{b,a}^s(z) = -\varphi_{-b,-a}^s(z) = \varphi_{-a,-b}^s(z).$$

Lemma 7.4. *For all $a, b \in \frac{1}{2} + \mathbb{Z}$ and $s \in \{0, 1\}$,*

$$\begin{aligned} \pi(\varphi_{a,b}^s(z^3 + z + z^{-1} + z^{-3})) &= \varphi_{a-1,b}^s + \varphi_{a+1,b}^s + \varphi_{a,b+1}^s + \varphi_{a,b-1}^s \\ \pi(\varphi_{a,b}^s(z^4 + z^2 + 1 + z^{-2} + z^{-4})) &= \varphi_{a+1,b+1}^s + \varphi_{a-1,b+1}^s + \varphi_{a+1,b-1}^s + \varphi_{a-1,b-1}^s + \varphi_{a,b}^s. \end{aligned}$$

(the projection π is introduced in Section 3).

Proof. Both equalities are straightforward corollaries of Lemma 7.2 and Lemma 3.2 (b) if one takes into account the isomorphisms of \mathfrak{k} -modules $V_{1,0} \simeq V_3$ and $V_{1,1} \simeq V_4$. \square

We define now $\psi_{a,b}^s(z) \in \mathbb{C}((z))$ via the conditions:

$$\begin{aligned} (c1) \quad \psi_{a,b}^s(z)(z^3 + z + z^{-1} + z^{-3}) &= \psi_{a+1,b}^s(z) + \psi_{a-1,b}^s(z) + \psi_{a,b+1}^s(z) + \\ &\quad \psi_{a,b-1}^s(z), \\ (c2) \quad \psi_{a,b}^s(z)(z^4 + z^2 + 1 + z^{-2} + z^{-4}) &= \psi_{a+1,b+1}^s(z) + \psi_{a+1,b-1}^s(z) + \\ &\quad \psi_{a-1,b+1}^s(z) + \psi_{a-1,b-1}^s(z) + \psi_{a,b}^s(z), \end{aligned}$$

$$(c3) \quad \psi_{a,b}^s(z) = -\psi_{b,a}^s(z) = -\psi_{-b,-a}^s(z) = \psi_{-a,-b}^s(z),$$

$$(c4) \quad \psi_{\frac{3}{2},\frac{1}{2}}^s(z) = \frac{z^s}{1-z^6}, \quad \psi_{\frac{3}{2},-\frac{1}{2}}^s(z) = \frac{z^{3+s}}{1-z^6}.$$

Theorem 7.5. *The Laurent series $\psi_{a,b}^s(z)$ exists and is unique. Moreover,*

(33)

$$\psi_{a,b}^s(z) = \frac{z^{5+s}(z^{3a+b} - z^{a+3b} - z^{-a-3b} + z^{-3a-b}) - z^{6+s}(z^{3a-b} - z^{-a+3b} - z^{a-3b} + z^{-3a+b})}{(1-z^2)^2(1-z^4)(1-z^6)}.$$

Proof. We show first that $\psi_{a,b}^s(z)$ is unique if it exists. By (32) $\psi_{a,b}^s(z)$ is determined by $\psi_{a,b}^s(z)$ for $a > |b|$. Assume, by induction on a , that $\psi_{a,b}^s(z)$ is unique for all $a \leq a_0$, $|b| < a$. Then equation (c1) determines $\psi_{a_0+1,b}^s(z)$, and equation (c2) determines $\psi_{a_0+1,a_0}^s(z)$ and $\psi_{a_0+1,a_0+1}^s(z)$.

To prove the existence of $\psi_{a,b}^s(z)$, it suffices to verify that the right-hand side of (33) satisfies all conditions (c1)-(c4). This is a direct calculation, which is simplified by the observation that both Laurent polynomials

$$\begin{aligned} & z^{3a+b} - z^{a+3b} - z^{-a-3b} + z^{-3a-b}, \\ & z^{3a-b} - z^{-a+3b} - z^{a-3b} + z^{-3a+b} \end{aligned}$$

satisfy (c1),(c2) and (c3). The condition (c4) is satisfied only by the entire expression. \square

Corollary 7.6.

$$\varphi_{a,b}^s = \pi(\psi_{a,b}^s).$$

Corollary 7.7. *Any simple bounded $(\mathfrak{g}, \mathfrak{k})$ -module is either even or odd. More precisely, $M_{a,b}^s$ is even if $a + b + s$ is even, and $M_{a,b}^s$ is odd if $a + b + s$ is odd.*

In the calculations below we use binomial coefficients $\binom{s}{k}$, for which we always assume $\binom{s}{k} = 0$ if s or k are not integers.

Lemma 7.8.

$$\frac{1}{(1-z^2)^2(1-z^4)(1-z^6)} = \sum_{n=0}^{\infty} \gamma(n)z^{2n},$$

where

$$\gamma(n) := \frac{1}{144} \left[119 \binom{n+3}{3} - 179 \binom{n+2}{3} + 109 \binom{n+1}{3} - 25 \binom{n}{3} \right] + \frac{(-1)^n}{16} + \frac{\beta(n)}{9}$$

and

$$\beta(n) := \begin{cases} 0 & n \equiv 1 \pmod{3} \\ 1 & n \equiv 0 \pmod{3} \\ -1 & n \equiv -1 \pmod{3} \end{cases}.$$

Proof. The statement follows from the identity

$$\frac{1}{(1-z^2)^2(1-z^4)(1-z^6)} = \frac{119-179z^2+109z^4-25z^6}{144(1-z^2)^4} + \frac{1}{16(1+z^2)} + \frac{1+z^2}{9(1+z^2+z^4)}.$$

□

Corollary 7.9. *Let*

$$\begin{aligned} \delta_{a,b}^s(n) = & \gamma\left(\frac{n-(3a+b+5)-s}{2}\right) - \gamma\left(\frac{n-(a+3b+5)-s}{2}\right) - \\ & -\gamma\left(\frac{n-(-a-3b+5)-s}{2}\right) + \gamma\left(\frac{n-(-3a-b+5)-s}{2}\right) - \\ & -\gamma\left(\frac{n-(3a-b+6)-s}{2}\right) + \gamma\left(\frac{n-(-a+3b+6)-s}{2}\right) + \\ & +\gamma\left(\frac{n-(a-3b+6)-s}{2}\right) - \gamma\left(\frac{n-(-3a+b+6)-s}{2}\right). \end{aligned}$$

Then

$$c_i(M_{a,b}^s) = \delta_{a,b}^s(i) - \delta_{a,b}^s(-i-2).$$

Proof. The statement follows directly from Theorem 7.5, Corollary 7.6, and Lemma 7.8. □

Corollary 7.10. *For any simple bounded $(\mathfrak{g}, \mathfrak{k})$ -module M , $c_i(M) = c_{i+6}(M)$ for sufficiently large $i \in \mathbb{N}$.*

Proof. The given $(\mathfrak{g}, \mathfrak{k})$ -module M is isomorphic to $M_{a,b}^s$ for some $a, b \in \frac{1}{2} + \mathbb{Z}$, $s \in \{0, 1\}$. For sufficiently large i , $\delta_{a,b}^s(-i-2) = 0$, hence $c_i(M) = \delta_{a,b}^s(i)$. The explicit formula for $\gamma(i)$ from Lemma 7.8 implies that $\delta_{a,b}^s(i+6n)$ is a polynomial in n . Since this polynomial is a bounded function, it is necessarily a constant. □

For large enough values of i , Corollary 7.10 enables us to write $c_{\bar{i}}(M_{a,b}^s)$, $\bar{i} \in \mathbb{Z}_6$. Here are simple explicit expressions for $c_{\bar{i}}(M_{a,b}^s)$.

Theorem 7.11. *Let $\sigma_{a,b} := \begin{cases} 1 & \text{if } 3|2a, 3 \nmid 2b \\ -1 & \text{if } 3|2b, 3 \nmid 2a \\ 0 & \text{in all other cases} \end{cases}$.*

Then

$$\begin{aligned} c_{\overline{0+s}}(M_{a,b}^s) &= \frac{1}{6}(1 + (-1)^{a+b}) \left(\frac{a^2 - b^2}{2} + 2\sigma_{a,b} \right), \\ c_{\overline{1+s}}(M_{a,b}^s) &= c_{\overline{5+s}}(M_{a,b}^s) = \frac{1}{6}(1 - (-1)^{a+b}) \left(\frac{a^2 - b^2}{2} - \sigma_{a,b} \right), \\ c_{\overline{2+s}}(M_{a,b}^s) &= c_{\overline{4+s}}(M_{a,b}^0) = \frac{1}{6}(1 + (-1)^{a+b}) \left(\frac{a^2 - b^2}{2} - \sigma_{a,b} \right), \end{aligned}$$

$$c_{\overline{3+s}}(M_{a,b}^s) = \frac{1}{6}(1 - (-1)^{a+b}) \left(\frac{a^2 - b^2}{2} + 2\sigma_{a,b} \right).$$

Proof. Let $\{\xi_{\bar{i}}\}_{\bar{i} \in \mathbb{Z}_6}$ denote the standard basis in \mathbb{C}^6 . Set

$$\overline{\varphi}_{a,b}^s := \sum_{\bar{i} \in \mathbb{Z}_6} c_{\bar{i}}(M_{a,b}^s) \xi_{\bar{i}}$$

for $a, b \in \frac{1}{2} + \mathbb{Z}$, $a \geq |b|$. Extend $\overline{\varphi}_{a,b}^s$ to all $a, b \in \frac{1}{2} + \mathbb{Z}$ by putting

$$\overline{\varphi}_{a,b}^s = -\overline{\varphi}_{b,a}^s = -\overline{\varphi}_{-b,-a}^s = \overline{\varphi}_{-a,-b}^s,$$

and let $S, T : \mathbb{C}^6 \rightarrow \mathbb{C}^6$ be the linear operators

$$S(\xi_{\bar{i}}) := 2\xi_{\overline{i+3}} + \xi_{\overline{i+1}} + \xi_{\overline{i-1}}, \quad T(\xi_{\bar{i}}) := 2\xi_{\overline{i+2}} + 2\xi_{\overline{i+4}}.$$

Then $\overline{\varphi}_{a,b}^s$ satisfy the following version of conditions (c1)-(c4):

- (c5) $S(\overline{\varphi}_{a,b}^s) = \overline{\varphi}_{a+1,b}^s + \overline{\varphi}_{a,b+1}^s + \overline{\varphi}_{a-1,b}^s + \overline{\varphi}_{a,b-1}^s$,
- (c6) $T(\overline{\varphi}_{a,b}^s) = \overline{\varphi}_{a+1,b+1}^s + \overline{\varphi}_{a-1,b+1}^s + \overline{\varphi}_{a+1,b-1}^s + \overline{\varphi}_{a-1,b-1}^s$,
- (c7) $\overline{\varphi}_{a,b}^s = -\overline{\varphi}_{b,a}^s = -\overline{\varphi}_{-b,-a}^s = \overline{\varphi}_{-a,-b}^s$,
- (c8) $\overline{\varphi}_{\frac{3}{2},\frac{1}{2}}^s = \xi_{\overline{s}}$, $\overline{\varphi}_{\frac{3}{2},-\frac{1}{2}}^s = \xi_{\overline{3+s}}$.

Denote by ω a primitive sixth root of unity. Then $\{\eta_{\bar{i}} := \sum_{\bar{j} \in \mathbb{Z}_6} \omega^{i\bar{j}} \xi_{\bar{j}}\}_{\bar{i} \in \mathbb{Z}_6}$

is an eigenbasis for S and T . Put

$$\begin{aligned} \eta_{\overline{0},a,b} &:= \frac{(a^2 - b^2)}{2} \eta_{\overline{0}}, & \eta_{\overline{3},a,b} &:= (-1)^{a+b} \frac{(a^2 - b^2)}{2} \eta_{\overline{3}}, \\ \eta_{\overline{2},a,b} &:= \sigma_{a,b} \eta_{\overline{2}}, & \eta_{\overline{4},a,b} &:= \sigma_{a,b} \eta_{\overline{4}}, \\ \eta_{\overline{3},a,b} &:= (-1)^{a+b} \sigma_{a,b} \eta_{\overline{3}}, & \eta_{\overline{5},a,b} &:= (-1)^{a+b} \sigma_{a,b} \eta_{\overline{5}}. \end{aligned}$$

Using the identity

$$\sigma_{a,b} = \frac{\omega^{2b} + \omega^{-2b} - \omega^{2a} - \omega^{-2a}}{3},$$

one can easily check that $\eta_{\bar{i},a,b}$ satisfies (c5)-(c7). The linear combination

$$\overline{\varphi}_{a,b}^s = \frac{1}{6} \sum_{\bar{i} \in \mathbb{Z}_6} \omega^{-i\bar{s}} \eta_{\bar{i},a,b}$$

satisfies the condition (c8), hence its coefficients in the basis $\{\xi_{\bar{i}}\}$ equal $c_{\bar{i}}(M_{a,b}^s)$. \square

Corollary 7.12. *The following is a complete list of multiplicity-free simple $(\mathfrak{g}, \mathfrak{k})$ -modules: $M_{\frac{3}{2},\pm\frac{1}{2}}^s, M_{\frac{5}{2},\pm\frac{3}{2}}^s, M_{\frac{5}{2},\pm\frac{1}{2}}^s, M_{\frac{7}{2},\pm\frac{5}{2}}^s$, $s \in \{0, 1\}$.*

Proof. A straightforward computation based on Theorem 7.11 shows that $c_{\bar{i}}(M_{a,b}^s) \in \{0, 1\}$ for $\bar{i} \in \mathbb{Z}_6$ iff (a, b) is one of the pairs $\left(\frac{3}{2}, \pm\frac{1}{2}\right)$, $\left(\frac{5}{2}, \pm\frac{3}{2}\right)$, $\left(\frac{5}{2}, \pm\frac{1}{2}\right)$, and $\left(\frac{7}{2}, \pm\frac{5}{2}\right)$. Then, using Corollary 7.9 one verifies that all modules $M_{a,b}^s$ for (a, b) as above are indeed multiplicity-free. \square

Theorem 7.13.

- (a) *The minimal \mathfrak{k} -type of any even (respectively, odd) bounded simple $(\mathfrak{g}, \mathfrak{k})$ -module M equals V_0, V_2 or V_4 (resp., V_1 or V_3).*
- (b) *If M is an even (respectively, odd) simple module in $\mathfrak{B}^{\chi(a,b)}$, then $c_0(M)$ (resp., $c_1(M)$) equals $\frac{a \pm b}{6} + \epsilon$ or $\frac{a \pm b}{12} + \epsilon$ (resp., $\frac{a \pm b}{3} + \epsilon$ or $\frac{a \pm b}{6} + \epsilon$) for some ϵ with $|\epsilon| < 1$.*

Proof. (a) Note that for any bounded $(\mathfrak{g}, \mathfrak{k})$ -module M , $c_i(M)$ equals the constant term of the Laurent polynomial $z^{-i}(1 - z^{2i+2})c(M)$. Hence $c_1(M) + c_3(M)$ equals the constant term in the Laurent expansion of $(z^{-1}(1 - z^4) + z^{-3}(1 - z^8))c(M)$. A straightforward calculation shows that for $M = M_{a,b}^s$ the latter is nothing but the constant term of the Laurent series

$$\frac{z^{3a+b+2+s} - z^{a+3b+2+s} - z^{-a-3b+2+s} + z^{-3a-b+2+s} - z^{-3a+b+3+s}}{(1 - z^2)^3} + \frac{z^{a-3b+3+s} + z^{-a+3b+3+s} - z^{3a-b+3+s}}{(1 - z^2)^3}.$$

Using the identity

$$(34) \quad \frac{1}{(1 - z^2)^3} = \sum_{n=0}^{\infty} \binom{n+2}{2} z^{2n},$$

we obtain

$$(35) \quad c_1(M_{a,b}^s) + c_3(M_{a,b}^s) =: d_{a,b}^s = \binom{\frac{-3a-b+2-s}{2}}{2} - \binom{\frac{-a-3b+2-s}{2}}{2} - \binom{\frac{a+3b+2-s}{2}}{2} + \binom{\frac{3a+b+2-s}{2}}{2} - \binom{\frac{3a-b+1-s}{2}}{2} + \binom{\frac{-a+3b+1-s}{2}}{2} + \binom{\frac{a-3b+1-s}{2}}{2} - \binom{\frac{-3a+b+1-s}{2}}{2},$$

where we set $\binom{l}{2} := 0$ for $l \notin \mathbb{Z}_{\geq 0}$.

This expression is a piecewise polynomial function which equals identically zero whenever $M_{a,b}^s$ is even, i.e. when $a + b + s$ is even. In fact,

the right hand side of (35) turns out to be very simple as an explicit calculation shows that, for $a + b + s$ odd,

$$(36) \quad d_{a,b}^s = \begin{cases} \frac{a + (-1)^{s+1}b}{2} & \text{for } a + (-1)^s 3b \geq 0 \\ a + (-1)^s b & \text{for } a + (-1)^s 3b \leq 0 \end{cases}.$$

Since $a > |b|$, the right hand side of (36) is never 0, i.e. the minimal \mathfrak{k} -type of $M_{a,b}^s$ is V_1 or V_3 whenever $a + b + s$ is odd.

A similar analysis proves that the minimal \mathfrak{k} -type of $M_{a,b}^s$ is V_0 , V_2 , or V_4 whenever $a + b + s$ is even. Indeed, in this case

$$e_{a,b}^s := c_0(M_{a,b}^s) + c_2(M_{a,b}^s) + c_4(M_{a,b}^s)$$

equals the constant term of the Laurent series

$$(1 - z^2) + z^{-2}(1 - z^6) + z^{-4}(1 - z^{10})c(M) \quad .$$

Using the identity

$$\frac{(1 - z^2) + z^{-2}(1 - z^6) + z^{-4}(1 - z^{10})}{(1 - z^2)^2(1 - z^4)(1 - z^6)} = \frac{1}{8z^4} \left(\frac{7 + 4z^2 + z^4}{(1 - z^2)^3} + \frac{1}{(1 + z^2)} \right),$$

as well as the identity (34), we calculate

$$\begin{aligned} e_{a,b}^s &= \theta \left(\frac{-3a - b - 1 - s}{2} \right) - \theta \left(\frac{-a - 3b - 1 - s}{2} \right) - \\ &\quad - \theta \left(\frac{a + 3b - 1 - s}{2} \right) + \theta \left(\frac{3a + b - 1 - s}{2} \right) - \\ &\quad - \theta \left(\frac{3a - b - 2 - s}{2} \right) + \theta \left(\frac{-a + 3b - 2 - s}{2} \right) + \\ &\quad + \theta \left(\frac{a - 3b - 2 - s}{2} \right) - \theta \left(\frac{-3a + b - 2 - s}{2} \right), \end{aligned}$$

where $\theta(n) := \frac{3}{4}n^2 + \frac{3}{2}n + \frac{7}{8} + \frac{(-1)^n}{8}$ for $n \in \mathbb{Z}_{\geq 0}$ and $\theta(n) := 0$ otherwise. Further calculations show:

$$(37) \quad e_{a,b}^s = \begin{cases} \frac{3}{4}(a + (-1)^{s+1}b) + \frac{(-1)^{\frac{a+(-1)^{s+1}b-1}{2}}}{4} & \text{for } (-1)^s a + 3b \geq 0 \\ \frac{3}{2}(a + (-1)^s b) & \text{for } (-1)^s a + 3b \leq 0 \end{cases}$$

under the assumption that $a + b + s$ is even. Since the right-hand side of (37) never equals 0, we obtain that $e_{a,b}^s \neq 0$ under the same assumption. Hence the minimal \mathfrak{k} -type of any even simple bounded $(\mathfrak{g}, \mathfrak{k})$ -module equals V_0 , V_2 , or V_4 .

(b) To compute $c_0(M)$ we use the identity

$$\begin{aligned} \frac{1-z^2}{(1-z^2)^2(1-z^4)(1-z^6)} &= \frac{1}{(1-z^2)(1-z^4)(1-z^6)} \\ &= \frac{47-52z^2+17z^4}{72(1-z^2)^3} + \frac{1}{8(1+z^2)} + \frac{2-z^2-z^4}{9(1-z^6)} \end{aligned}$$

which yields

$$\begin{aligned} c_0(M_{a,b}^s) &= \gamma' \left(\frac{-3a-b-5-s}{2} \right) - \gamma' \left(\frac{-a-3b-5-s}{2} \right) - \\ &\quad - \gamma' \left(\frac{a+3b-5-s}{2} \right) + \gamma' \left(\frac{3a+b-5-s}{2} \right) - \\ &\quad - \gamma' \left(\frac{3a-b-6-s}{2} \right) + \gamma' \left(\frac{-a+3b-6-s}{2} \right) + \\ &\quad + \gamma' \left(\frac{a-3b-6-s}{2} \right) - \gamma' \left(\frac{-3a+b-6-s}{2} \right), \end{aligned}$$

where

$$\begin{aligned} \gamma'(n) &:= \frac{n^2}{12} + \frac{n}{2} + \frac{94}{144} + \frac{(-1)^n}{8} + \frac{\sigma'(n)}{9}, \\ \sigma'(n) &:= \begin{cases} -1 & 3 \nmid n \\ 2 & 3 \mid n \end{cases} \end{aligned}$$

for $n \in \mathbb{Z}_{\geq 0}$ and $\gamma'(n) = \sigma'(n) := 0$ otherwise. Similarly, using the identity

$$\frac{z^{-1}(1-z^4)}{(1-z^2)^2(1-z^4)(1-z^6)} = z^{-1} \left(\frac{8-7z^2+2z^4}{9(1-z^2)^3} + \frac{1+z^2-2z^4}{9(1-z^6)} \right)$$

we obtain

$$\begin{aligned} c_1(M_{a,b}^s) &= \gamma'' \left(\frac{-3a-b-4-s}{2} \right) - \gamma'' \left(\frac{-a-3b-4-s}{2} \right) - \\ &\quad - \gamma'' \left(\frac{a+3b-4-s}{2} \right) + \gamma'' \left(\frac{3a+b-4-s}{2} \right) - \\ &\quad - \gamma'' \left(\frac{3a-b-5-s}{2} \right) + \gamma'' \left(\frac{-a+3b-5-s}{2} \right) + \\ &\quad + \gamma'' \left(\frac{a-3b-5-s}{2} \right) - \gamma'' \left(\frac{-3a+b-5-s}{2} \right), \end{aligned}$$

where

$$\gamma''(n) := \frac{n^2}{6} + \frac{5n}{6} + \frac{8}{9} + \frac{\sigma''(n)}{9},$$

$$\sigma''(n) := \begin{cases} -2 & n = -1 \pmod{3} \\ 1 & n \neq -1 \pmod{3} \end{cases}$$

for $n \in \mathbb{Z}_{\geq 0}$ and $\gamma''(n) = \sigma''(n) := 0$ otherwise. Using the expressions for $c_0(M_{a,b}^s)$ and $c_1(M_{a,b}^s)$ we notice that the terms $\frac{(-1)^n}{8} + \frac{\sigma'(n)}{9}$ and $\frac{\sigma''(n)}{9}$ will give a contribution ϵ with $|\epsilon| < 1$. Thus, a direct computation implies

$$c_0(M_{a,b}^s) = \begin{cases} \frac{a+(-1)^s b}{6} + \epsilon & \text{for } a + (-1)^s 3b < 0 \\ \frac{a-(-1)^s b}{12} + \epsilon & \text{for } a + (-1)^s 3b > 0 \end{cases},$$

$$c_1(M_{a,b}^s) = \begin{cases} \frac{a-(-1)^s b}{6} + \epsilon & \text{for } a + (-1)^s 3b > 0 \\ \frac{a+(-1)^s b}{3} + \epsilon & \text{for } a + (-1)^s 3b < 0. \end{cases}$$

□

Corollary 7.14. *For $a \pm b \geq 24$, the minimal \mathfrak{k} -type of $M_{a,b}^s$ equals V_0 (respectively, V_1) if $a + b + s$ is odd (resp., even).*

Corollary 7.15. *A simple $(\mathfrak{g}, \mathfrak{k})$ -module with minimal \mathfrak{k} -type V_i for $i \geq 5$ is unbounded.*

Note that all simple $(\mathfrak{g}, \mathfrak{k})$ -modules of finite type over \mathfrak{k} with minimal \mathfrak{k} -type V_i for $i \geq 6$ are classified in [PZ2]. In particular it is proved, [PZ2], that if M is a $(\mathfrak{g}, \mathfrak{k})$ -module with minimal \mathfrak{k} -type V_i for $i \geq 6$, then M is necessarily of finite type over \mathfrak{k} and $c_i(M) = 1$. Recently G. Zuckerman and the first named author have shown that this holds also for $i = 5$, and Theorem 7.13 (b) implies that the statement is false for $i \leq 1$.

REFERENCES

- B. V. Bargmann, Irreducible unitary representations of the Lorentz group, *Ann. of Math. (2)* **48** (1947), 586–640.
- BG. J. Bernstein, S. Gelfand, Tensor products of finite and infinite-dimensional representations of semi-simple Lie algebras, *Compositio Math.* **41** (1980), 245–285.
- Dix. J. Dixmier, *Enveloping algebras*, North-Nolland Math. Library, 14, North Holland, Amsterdam, 1977.
- F. S. L. Fernando, Lie algebra modules with finite-dimensional weight spaces I, *Trans. Amer. Math. Soc.* **322** (1990), 757–781.
- Fo. A. I. Fomin, Quasisimple irreducible representations of the group $SL(3, \mathbb{R})$. *Func. Anal. i Pril.* **9** (1975), No 3, 67–74.
- GN. I. M. Gelfand, M. A. Naimark, Unitary representations of the Lorentz group, *Izvestiya Akad. Nauk SSSR, Ser. Mat.* **11** (1947), 411–504.
- HC. Harish-Chandra, Infinite irreducible representations of the Lorentz group, *Proc. Roy. Soc. London, Ser. A* **189** (1947), 372–401.

- Kn1. F. Knop, Der Zentralisator einer Liealgebra in einer einhüllenden Algebra (German), *J. Reine Angew. Math.* **406** (1990), 5–9.
- Kn2. F. Knop, Classification of multiplicity-free symplectic representations. *J. Algebra* **301** (2006), 531–553.
- KV. A. W. Knap, D. Vogan, Jr., Cohomological induction and unitary representations, Princeton University Press, Princeton, NJ, 1995.
- OV. A. Onishchik, E. Vinberg, Lie groups and algebraic groups. Springer series in Soviet mathematics, Springer-Verlag, Berlin, 1990.
- PS1. I. Penkov, V. Serganova, Generalized Harish-Chandra modules, *Moscow Math. J.* **2** (2002), 753–767.
- PS2. I. Penkov, V. Serganova, On bounded generalized Harish-Chandra modules, preprint, 2009, math.jacobs-university.de/penkov
- PSZ. I. Penkov, V. Serganova, G. Zuckerman, On the existence of $(\mathfrak{g}, \mathfrak{k})$ -modules of finite type, *Duke Math. J.* **125** (2004), 329–349.
- PZ1. I. Penkov, G. Zuckerman, Generalized Harish-Chandra modules: a new direction of the structure theory of representations, *Acta Applic. Math.* **81** (2004), 311–326.
- PZ2. I. Penkov, G. Zuckerman, Generalized Harish-Chandra modules with generic minimal \mathfrak{k} -type, *Asian J. of Math.* **8** (2004), 795–812.
- PZ3. I. Penkov, G. Zuckerman, A construction of generalized Harish-Chandra modules with arbitrary minimal \mathfrak{k} -type, *Canad. Math. Bull.* **50** (2007), 603–609.
- S. Dj. Sijacki, All $SL(3, \mathbb{R})$ ladder representations, *J. Math. Phys.* **31** (1990), 1872–1876.
- Sp. B. Speh, The unitary dual of $GL(3, \mathbb{R})$ and $GL(4, \mathbb{R})$, *Ann. Math.* **258**(1981/82), 113–133.
- V1. D. Vogan, Jr., Singular unitary representations. In *Lecture Notes in Mathematics* 880, Springer-Verlag, 1981.
- V2. D. Vogan, Jr., The unitary dual of G_2 . *Invent. Math.* **116** (1994), 677–791.

JACOBS UNIVERSITY BREMEN, SCHOOL OF ENGINEERING AND SCIENCE, CAMPUS RING 1, 28759 BREMEN, GERMANY

E-mail address: i.penkov@jacobs-university.de

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA BERKELEY, BERKELEY CA 94720, USA

E-mail address: serganov@math.berkeley.edu