## Week 13: More Matrices

1. MULTI Single Find the inverse of  $A = \begin{vmatrix} 0 & \frac{1}{2} & -\frac{1}{2} \\ 1 & 0 & 1 \\ 2 & \frac{1}{2} & 1 \end{vmatrix}$ . (a)  $A^{-1} = \begin{bmatrix} -2 & -3 & 2 \\ 4 & 4 & 2 \\ 2 & 4 & -2 \end{bmatrix}$ (b)  $A^{-1} = \begin{bmatrix} -2 & -3 & 2 \\ -2 & -3 & 2 \\ 4 & 4 & 2 \\ 2 & -4 & -2 \end{bmatrix}$ (c) the inverse does not exist (d)  $A^{-1} = \begin{bmatrix} -2 & -3 & 2 \\ 4 & 4 & -2 \\ 2 & 4 & -2 \end{bmatrix}$ 2.MULTI Single Find the inverse of  $A = \begin{bmatrix} 3.5 & -1 & 0.5 \\ 10 & -3 & 2 \\ 2.5 & -1 & 1.5 \end{bmatrix}$ . (a)  $A^{-1} = \begin{bmatrix} 0 & 1 & 1 \\ 2 & -1 & -2 \\ -1 & 1 & -2 \end{bmatrix}$ (b)  $A^{-1} = \begin{bmatrix} 0 & 1 & 1 \\ 2 & -1 & -2 \\ 1 & 1 & 2 \end{bmatrix}$ (c) the inverse does not exist (d)  $A^{-1} = \begin{bmatrix} 0 & 1 & 1 \\ 2 & -1 & -2 \\ 1 & 1 & -2 \end{bmatrix}$ 3. MULTI Single Find the inverse of  $A = \begin{vmatrix} 1 & -1 & 1 \\ -1 & 1 & 1 \\ 1 & 1 & 1 \end{vmatrix}$ . (a)  $A^{-1} = \begin{bmatrix} 0.5 & -0.5 & 0.5 \\ -0.5 & 0 & 0.5 \\ 0.5 & -0.5 & 0 \end{bmatrix}$ (b)  $A^{-1} = \begin{bmatrix} 0 & -0.5 & 0.5 \\ -0.5 & 0 & 0.5 \\ 0.5 & 0.5 & 0 \end{bmatrix}$ (c)  $A^{-1} = \begin{bmatrix} 0.5 & -0.5 & 0 \\ -0.5 & 0 & 0.5 \\ 0.5 & 0.5 & 0 \end{bmatrix}$ (d) the inverse does not exist

4. MULTI Single

Find the kernel of 
$$A = \begin{bmatrix} 1 & 2 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$
.  
(a)  $\ker(A) = \operatorname{span} \left\{ \begin{bmatrix} 2 \\ 1 \\ -1 \end{bmatrix} \right\}$   
(b)  $\ker(A) = \operatorname{span} \left\{ \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix} \right\}$   
(c)  $\ker(A) = \operatorname{span} \left\{ \begin{bmatrix} 2 \\ 1 \\ 0 \end{bmatrix} \right\}$   
(d)  $\ker(A) = \operatorname{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ -1 \end{bmatrix} \right\}$ 

5. MULTI Single

An  $n \times k$  matrix A has the following kernel:

$$\ker(A) = \operatorname{span}\left\{ \begin{bmatrix} 2\\1\\0 \end{bmatrix}, \begin{bmatrix} -2\\1\\0 \end{bmatrix} \right\}$$

What is the dimension of its image  $\dim(\operatorname{im}(A))$ ?

- (a)  $\dim(\operatorname{im}(A)) = n 3$ (b)  $\dim(\operatorname{im}(A)) = k - 1$ (c)  $\dim(\operatorname{im}(A)) = k - 2$ (d)  $\dim(\operatorname{im}(A)) = n$
- 6. MULTI Single

Given the matrix  $A = B \cdot C \cdot D$  find its inverse  $A^{-1}$ .

- $\begin{array}{ll} ({\rm a}) & A^{-1} = C^{-1}D^{-1}B^{-1}\\ ({\rm b}) & A^{-1} = D^{-1}B^{-1}C^{-1}\\ ({\rm c}) & A^{-1} = D^{-1}C^{-1}B^{-1}\\ ({\rm d}) & A^{-1} = B^{-1}C^{-1}D^{-1} \end{array}$
- 7. Multi Single

A linear map  $D : \mathbb{R}^2 \to \mathbb{R}^2$  is given by the following matrix in the standard basis:  $D_{\text{st}} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ . How is the map represented in the following basis:  $\left\{ \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \frac{1}{\sqrt{2}} \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$ ? (a)  $D_{\text{new}} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \\ 0 & 1 \end{bmatrix}$ (b)  $D_{\text{new}} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ 

- (c)  $D_{\text{new}} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ (d)  $D_{\text{new}} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$
- 8. MULTI Single

Consider the standard basis in  $\mathbb{R}^3$ :  $\{e_x, e_y, e_z\}$ .

Which of the following matrices represents a counterclockwise rotation around the z-axis?

(a) 
$$\mathcal{R} = \begin{bmatrix} \cos \varphi & 0 & \sin \varphi \\ 0 & 1 & 0 \\ \sin \varphi & 0 & \cos \varphi \end{bmatrix}$$
  
(b) 
$$\mathcal{R} = \begin{bmatrix} 1 & 0 & -\sin \varphi \\ \cos \varphi & 1 & \cos \varphi \\ \sin \varphi & 0 & 1 \end{bmatrix}$$
  
(c) 
$$\mathcal{R} = \begin{bmatrix} \cos \varphi & -\sin \varphi & 0 \\ \sin \varphi & \cos \varphi & 0 \\ 0 & 0 & 1 \end{bmatrix}$$
  
(d) 
$$\mathcal{R} = \begin{bmatrix} \cos \varphi & -\sin \varphi & 0 \\ \sin \varphi & \cos \varphi & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

9. MULTI Single

Consider the space of  $2 \times 2$  Hermitian Matrices  $H_2(\mathbb{C})$  (the space of  $2 \times 2$  matrices A with complex entries such that  $A^{\dagger} = A$ ). Which of the following is true?

- (a)  $H_2(\mathbb{C})$  is a vector space over the field of complex numbers, but not over the real numbers.
- (b)  $H_2(\mathbb{C})$  is not a vector space over the field of real numbers or complex numbers.
- (c)  $H_2(\mathbb{C})$  is a vector space over the field of real numbers, but not over the complex numbers.
- (d)  $H_2(\mathbb{C})$  is a vector space over the field of real numbers and over the complex numbers.
- 10. MULTI Single

Consider the vector space  $P_2(\mathbb{R}) = \{p(x) \mid p(x) \text{ is a quadratic polynomial}\}.$ Is the derivative operator  $\mathcal{D} : p(x) \mapsto p'(x) \equiv \frac{\mathrm{d}}{\mathrm{d}x}p(x)$  a linear operator? If it is, how is it represented in the standard basis  $\mathfrak{B} = \{1, x, x^2\}$ 

*Hint:* You can express a polynomial  $ax^2 + bx + c$  as  $\begin{bmatrix} c \\ b \\ a \end{bmatrix}$ 

- (a)  $\mathcal{D}$  is a linear operator with  $[\mathcal{D}]_{\mathfrak{B}} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 2 \\ 0 & 0 & 0 \end{bmatrix}$
- (b)  $\mathcal{D}$  is not a linear operator

(c) 
$$\mathcal{D}$$
 is a linear operator with  $[\mathcal{D}]_{\mathfrak{B}} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$   
(d)  $\mathcal{D}$  is a linear operator with  $[\mathcal{D}]_{\mathfrak{B}} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ 

Total of marks: 10