

2.6 Black-Scholes Formula

Recall the exact formula for the price of European call options:

$$C = e^{-rT} \sum_{j=0}^n \binom{n}{j} p^j (1-p)^{n-j} \max(0, S u^j d^{n-j} - K) \quad (K \text{ strike price, } r \text{ period interest rate})$$

$$= e^{-rT} \mathbb{E}(\text{payoff})$$

↳ under binomial distribution

Many terms in the sum are 0; payoff $\neq 0$ if $S u^j d^{n-j} - K > 0$

$$\Rightarrow \text{need } u^j d^{n-j} > \frac{K}{S} \Rightarrow \left(\frac{u}{d}\right)^j > \frac{K}{S d^n} \Rightarrow j > \frac{\ln\left(\frac{K}{S d^n}\right)}{\ln\left(\frac{u}{d}\right)} = \frac{\ln\left(\frac{K}{S}\right) - n \ln d}{\ln\left(\frac{u}{d}\right)} =: A_n$$

$$\Rightarrow C = e^{-rT} \sum_{j=A_n}^n \binom{n}{j} p^j (1-p)^{n-j} (S u^j d^{n-j} - K)$$

$$= S \sum_{j=A_n}^n \binom{n}{j} (p u e^{-r\frac{T}{n}})^j ((1-p)d e^{-r\frac{T}{n}})^{n-j} - K e^{-rT} \sum_{j=A_n}^n \binom{n}{j} p^j (1-p)^{n-j}$$

$$\text{Recall } p = \frac{e^{r\frac{T}{n}} - d}{u - d} \Rightarrow 1-p = \frac{u - d - e^{r\frac{T}{n}} + d}{u - d} = \frac{u - e^{r\frac{T}{n}}}{u - d}$$

$$\Rightarrow (1-p)d e^{-r\frac{T}{n}} = \frac{u - e^{r\frac{T}{n}}}{u - d} d e^{-r\frac{T}{n}} = \frac{u d e^{-r\frac{T}{n}} - d + u - u}{u - d} = 1 - p u e^{-r\frac{T}{n}}$$

$$\Rightarrow C = S \sum_{j=A_n}^n b(j, n, p u e^{-r\frac{T}{n}}) - K e^{-rT} \sum_{j=A_n}^n b(j, n, p)$$

Next: use our calibration $u = e^{6\sqrt{\frac{T}{n}}}$, $d = \frac{1}{u} = e^{-6\sqrt{\frac{T}{n}}}$

$$\Rightarrow A_n = \frac{\ln(\frac{k}{S}) - n \ln d}{\ln(\frac{k}{d})} = \frac{\ln(\frac{k}{S}) - n \ln e^{-6\sqrt{\frac{T}{n}}}}{\ln e^{26\sqrt{\frac{T}{n}}}} = \frac{\ln(\frac{k}{S}) + 6\sqrt{T}\sqrt{n}}{26\sqrt{\frac{T}{n}}} = \frac{\ln(\frac{k}{S})}{26\sqrt{T}}\sqrt{n} + \frac{1}{2}n$$

$$\Rightarrow p = \frac{e^{r\frac{T}{n}} - d}{u - d} = \frac{e^{r\frac{T}{n}} - e^{-6\sqrt{\frac{T}{n}}}}{e^{6\sqrt{\frac{T}{n}}} - e^{-6\sqrt{\frac{T}{n}}}} \stackrel{\text{Taylor}}{=} \frac{1 + r\frac{T}{n} + \underbrace{O(n^{-2})}_{\text{"big O" notation}} - \left[1 - 6\sqrt{\frac{T}{n}} + \frac{1}{2}6^2\frac{T}{n} + O(n^{-\frac{3}{2}}) \right]}{1 + 6\sqrt{\frac{T}{n}} + \frac{1}{2}6^2\frac{T}{n} + O(n^{-\frac{3}{2}}) - \left[1 - 6\sqrt{\frac{T}{n}} + \frac{1}{2}6^2\frac{T}{n} + O(n^{-\frac{3}{2}}) \right]}$$

$$= \frac{6\sqrt{\frac{T}{n}} + (r - \frac{6^2}{2})\frac{T}{n} + O(n^{-\frac{3}{2}})}{26\sqrt{\frac{T}{n}} + O(n^{-\frac{3}{2}})}$$

$$= \frac{1}{2} \left(1 + \frac{(r - \frac{6^2}{2})}{6} \sqrt{\frac{T}{n}} + O(n^{-1}) \right)$$

$$= \frac{O(n^{-\frac{3}{2}})}{26\sqrt{T}n^{-\frac{1}{2}}}$$

Then the integral boundary from the CLT is

$$\hat{A} = \lim_{n \rightarrow \infty} \frac{A_n - np}{\sqrt{np(1-p)}} = \lim_{n \rightarrow \infty} \frac{\frac{\ln(\frac{k}{S})}{26\sqrt{T}}\sqrt{n} + \frac{1}{2}n - \left[\frac{n}{2} + \frac{n}{2} \frac{(r - \frac{6^2}{2})}{6} \sqrt{\frac{T}{n}} + O(n^{-1}) \right]}{\sqrt{n} \left(\frac{1}{2} + \frac{\text{const}}{\sqrt{n}} + O(n^{-1}) \right)^{\frac{1}{2}} \left(\frac{1}{2} - \frac{\text{const}}{\sqrt{n}} + O(n^{-1}) \right)^{\frac{1}{2}}}$$

$$= \sqrt{n} \sqrt{\frac{1}{4} + O(n^{-1})} = \sqrt{n} \frac{1}{2} + O(n^{-\frac{1}{2}})$$

$$= \frac{\ln(\frac{k}{S}) - (r - \frac{6^2}{2})T}{6\sqrt{T}}$$

Now applying the CLT gives us: (after some more computation)

$$C = S \Phi(x) - Ke^{-rT} \Phi(x - \sigma\sqrt{T})$$
, the Black-Scholes formula,

with cumulative normal distribution $\Phi(x) = \int_{-\infty}^x \varphi(y) dy = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{y^2}{2}} dy$
normal gaussian

$$\text{and } x = \frac{\ln\left(\frac{S}{K}\right) + \left(r + \frac{\sigma^2}{2}\right)T}{\sigma\sqrt{T}}$$

let's check: $\hat{A} = \frac{\ln\left(\frac{K}{S}\right) - \left(r - \frac{\sigma^2}{2}\right)T}{\sigma\sqrt{T}} = -\left(\frac{\ln\left(\frac{S}{K}\right) + \left(r - \frac{\sigma^2}{2}\right)T}{\sigma\sqrt{T}}\right)$
 $= -(x - \sigma\sqrt{T})$

Then: $Ke^{-rT} \sum_{j=A_n}^n b(j, n, p) \xrightarrow[\text{CLT}]{n \rightarrow \infty} Ke^{-rT} \int_A^{\infty} \varphi(y) dy = Ke^{-rT} \int_{-(x - \sigma\sqrt{T})}^{\infty} \varphi(y) dy$
 $= Ke^{-rT} \int_{-\infty}^{x - \sigma\sqrt{T}} \varphi(y) dy$
 $= Ke^{-rT} \Phi(x - \sigma\sqrt{T})$

Similar for the first summand, but we need to use approximations for $pe^{-r\frac{T}{n}}$.

2.7 Convergence Rates

Consider some sequence $C_n \xrightarrow{n \rightarrow \infty} C$, e.g., $C_n =$ option price for n step binomial tree, and

$C =$ price from Black-Scholes

It is very important to know how fast the convergence is.

Often, the convergence goes like a power law: $E_n = |C_n - C| \approx A n^{-\beta}$ for large n


β is called rate of convergence

Remarks: • different β can make a huge difference (e.g. linear vs. quadratic)


• if limit C is unknown one could, e.g., look at $N \gg n$ and then

$$|C_n - C_N|, \text{ or } E_n = |C_{n+1} - C_n|$$

How can we read off β from a plot?



better: $\ln E_n = \ln A n^{-\beta} = \ln A - \beta \ln n$



python: $\log(\log(n, E_n))$ produces such plots