

Advanced Calculus and Methods of Mathematical Physics

Homework 4

Due on March 7, 2023, before the tutorial.

Problem 1 [4 points]

Consider the function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by

$$f(x, y) = \begin{cases} \frac{xy^2}{x^2 + y^2} & \text{when } (x, y) \neq (0, 0), \\ 0 & \text{when } (x, y) = (0, 0). \end{cases}$$

- (a) Compute the directional derivative $D_{\mathbf{u}}f|_{(0,0)}$ for every $\mathbf{u} = (u_1, u_2) \in \mathbb{R}^2$ with $\|\mathbf{u}\| = 1$. Is $\mathbf{u} \mapsto D_{\mathbf{u}}f|_{(0,0)}$ linear?
- (b) Show that f is continuous, but not differentiable at the origin.

Problem 2 [4 points]

Consider the function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by

$$f(x, y) = \begin{cases} \frac{x^3 y}{x^6 + y^2} & \text{when } (x, y) \neq (0, 0), \\ 0 & \text{when } (x, y) = (0, 0). \end{cases}$$

- (a) Compute the directional derivative $D_{\mathbf{u}}f|_{(0,0)}$ for every $\mathbf{u} = (u_1, u_2) \in \mathbb{R}^2$ with $\|\mathbf{u}\| = 1$. Is $\mathbf{u} \mapsto D_{\mathbf{u}}f|_{(0,0)}$ linear?
- (b) Show that f is not continuous at the origin.

Problem 3 [4 points]

Consider the function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by

$$f(x, y) = \begin{cases} \frac{x^2 y}{x^4 + y^2} \sqrt{x^2 + y^2} & \text{when } (x, y) \neq (0, 0), \\ 0 & \text{when } (x, y) = (0, 0). \end{cases}$$

- (a) Compute the directional derivative $D_{\mathbf{u}}f|_{(0,0)}$ for every $\mathbf{u} = (u_1, u_2) \in \mathbb{R}^2$ with $\|\mathbf{u}\| = 1$. Is $\mathbf{u} \mapsto D_{\mathbf{u}}f|_{(0,0)}$ linear?

(b) Show that f is continuous, but not differentiable at the origin.

Problem 4 [4 points]

Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ be differentiable on $\mathbb{R}^2 \setminus \{0\}$. Let

$$h(r, \theta) = (r \cos \theta, r \sin \theta)$$

denote the change from polar to Cartesian coordinates and set $g = f \circ h$. Prove that, for $r > 0$,

$$\|(\nabla f) \circ h\|^2 = \left(\frac{\partial g}{\partial r}\right)^2 + \left(\frac{1}{r} \frac{\partial g}{\partial \theta}\right)^2.$$

Problem 5 [4 points]

Let $\text{Mat}(n \times n)$ denote the set of real $n \times n$ matrices. We consider the squaring map

$$S: \text{Mat}(n \times n) \rightarrow \text{Mat}(n \times n), \quad S(A) \mapsto A^2.$$

In analogy to what we have defined in class, the map S is differentiable at A if there exists a linear map $DS|_A$ such that

$$S(A + H) = S(A) + DS|_A H + r_A(H), \quad \text{with } \lim_{H \rightarrow 0} \frac{\|r_A(H)\|}{\|H\|} = 0.$$

Here, $\|H\|$ denotes the operator norm of H . Show that A is differentiable everywhere and compute its derivative $DS|_A$.