

# Advanced Calculus and Methods of Mathematical Physics

## Homework 6

Due on March 21, 2023, before the tutorial.

### Problem 1 [3 points]

(Kantorovitz, p. 78, Exercise 6. Warm-up.) Let  $f: \mathbb{R}^k \rightarrow \mathbb{R}^m$  be defined by

$$f(x) = \sum_{i=1}^k (x_i, x_i^2, \dots, x_i^m).$$

Compute the derivative  $Df|_x$ .

### Problem 2 [6 points]

(From Rudin, Exercise 9.17.) Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be defined by

$$f(x) = \begin{pmatrix} e^{x_1} \cos x_2 \\ e^{x_1} \sin x_2 \end{pmatrix}.$$

- What is the range of  $f$ ?
- Show that the *Jacobian determinant*,  $\det Df|_x$ , is non-zero for every  $x \in \mathbb{R}^2$ . Thus every point in  $\mathbb{R}^2$  has a neighborhood in which  $f$  is one-to-one. Nevertheless,  $f$  is not one-to-one on  $\mathbb{R}^2$ .
- Put  $a = (0, \pi/3)$  and  $b = f(a)$ . Find an explicit formula for  $f^{-1}$ , compute  $Df|_a$  and  $Df^{-1}|_b$ , and verify the formula for the derivative of the inverse from the statement of the inverse function theorem (i.e.,  $Df^{-1}|_b = (Df|_a)^{-1}$ ).
- What are the images under  $f$  of lines parallel to the coordinate axes?

### Problem 3 [4 points]

(Kantorovitz, p. 106, Exercise 1.) Show that the equation

$$x^5 + y^5 + z^5 = 2 + xyz$$

determines in a neighborhood of the point  $(1, 1, 1)$  a unique function  $z = z(x, y)$  of class  $C^1$ , and calculate its partial derivatives with respect to  $x$  and  $y$  at the point  $(1, 1)$ .

**Problem 4 [3 points]**

Consider the equation

$$\sqrt{x^2 + y^2 + 2z^2} = \cos z$$

near  $(0, 1, 0)$ . Can you solve for  $x$  in terms of  $y$  and  $z$ ? For  $z$  in terms of  $x$  and  $y$ ?

**Problem 5 [4 points]**

Show that if  $r$  is a simple root of the polynomial

$$p(x) = a_0 + a_1 x + \cdots + a_n x^n,$$

then  $r$  is a  $C^1$  function of the coefficients  $a_0, \dots, a_n$ .