

Mock Exam
Calculus and Linear Algebra I
Fall 2022

Dr. Stephan Juricke, Prof. Soeren Petrat

1. **(6 points)** We consider the function

$$f(x) = e^{2x+1}.$$

Which of the following is true?

- A. The domain of f is $[0, \infty)$.
 - B. The domain of f is \mathbb{R} .
 - C. The domain of f is the set of all x with $x > -\frac{1}{2}$.
 - D. The inverse function is $f^{-1}(y) = \frac{1}{2} \ln(y) - \frac{1}{2}$.
 - E. The inverse function does not exist, since f is not invertible.
 - F. The inverse function is $f^{-1}(y) = e^{-2y-1}$.
2. **(6 points)** In class we discussed the Extreme Value Theorem: If $f : [a, b] \rightarrow \mathbb{R}$ is continuous, then f assumes its minimum and maximum. Which of the following is true?
- A. The theorem applies to $f : [0, 4] \rightarrow \mathbb{R}, f(x) = e^{\sin(x)}$.
 - B. The theorem applies to $f : (0, 1) \rightarrow \mathbb{R}, f(x) = \frac{1}{x}$.
 - C. The theorem applies to $f : [1, 2] \rightarrow \mathbb{R}, f(x) = \frac{1}{x}$.
 - D. The theorem applies to $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = e^x$.
 - E. The theorem applies to $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = \frac{x^2+1}{x^2+2}$.
 - F. The theorem applies to $f : [0, 1] \rightarrow \mathbb{R}, f(x) = \frac{x^2+1}{x^2+2}$.

3. (6 points) Consider the polynomial

$$p(x) = 2x^2 - 12x + 20.$$

Which of the following is true?

- A. The roots of p are $3 + i$ and $3 - i$.
- B. The roots of p are $3 + i$ and $3 + i$.
- C. The roots of p are $2 + i$ and $1 - 2i$.
- D. The roots of p are $1 + 3i$ and $1 - 3i$.
- E. $p(x)$ does not have any (possibly complex) roots.
- F. $p(x) > 0$ for all $x \in \mathbb{R}$.

4. (6 points) Consider the function

$$f(x) = \frac{\sin(x)}{x} + \frac{x}{x^2 - 1}.$$

Which of the following is true?

- A. The line $y = 0$ is a horizontal asymptote of f .
- B. The line $y = 1$ is a horizontal asymptote of f .
- C. f has a vertical asymptote at 0.
- D. f has a vertical asymptote at 1.
- E. f has a vertical asymptote at -1 .
- F. $f(0) = 0$.

5. **(6 points)** If the derivative f' of $f : [0, 1] \rightarrow \mathbb{R}$ at x_0 exists, which of the following statements is true?

- A. f is continuous at x_0 .
- B. f'' exists at x_0 .
- C. f is differentiable on $[0, 1]$.
- D. $f'(x_0) = \lim_{x \rightarrow x_0} \frac{f(x) - f(x_0)}{x - x_0}$.
- E. f' is continuous at x_0 .
- F. f is a polynomial.

6. **(4 points)** What is the derivative of $f(x) = \cos\left(\frac{\sqrt{e^x + a}}{2}\right)$ with respect to x , where $a \in \mathbb{R}$ is some constant?

- A. $f'(x) = e^x \frac{\cos\left(\frac{\sqrt{e^x + a}}{2}\right)}{\sqrt{e^x + a}}$
- B. $f'(x) = -e^x \sin\left(\frac{\sqrt{e^x + a}}{2}\right)$
- C. $f'(x) = -e^x \frac{\sin\left(\frac{\sqrt{e^x + a}}{2}\right)}{4\sqrt{e^x + a}}$
- D. $f'(x) = e^x \frac{\sin\left(\frac{\sqrt{e^x + a}}{2}\right)}{\sqrt{e^x + a}}$
- E. $f'(x) = -2e^x \frac{\sin\left(\frac{\sqrt{e^x + a}}{2}\right)}{x\sqrt{e^x + a}}$
- F. $f'(x) = -2e^x \frac{\tan\left(\frac{\sqrt{e^x + a}}{2}\right)}{x\sqrt{e^x + a}}$

7. (4 points) We need to build a box whose base length L is 3 times its base width W . The height is given by H . Different material is used to build the top and bottom. For the top it costs 10 EUR per square metre and for the sides it costs 6 EUR per square metre. The box should have a volume of 50 cubic metres. What is the value of the width of the base W that minimize the cost to build the box?

- A. $W = (3/20)^{1/3}$
- B. $W = (20/5)^{1/3}$
- C. $W = (20/9)^{1/3}$
- D. $W = (-9/20)^{1/3}$
- E. $W = (5/3)^{1/3}$
- F. $W = (20/3)^{1/3}$

8. (4 points) Compute the integral

$$\int_0^1 x^2 \sin(x^3 + 1) dx.$$

Hint: Substitution.

- A. $\frac{1}{3}(\cos(1) - 1)$.
- B. $\cos(2) - \cos(1)$.
- C. $\frac{1}{3}(\cos(1) - \cos(2))$.
- D. $\sin(2)$.
- E. $\sin(1)$.
- F. $\tan(2) - \tan(1)$.

9. (4 points) What is the solution to the differential equation

$$\frac{dy}{dt} = y + 1$$

with initial condition $y(0) = 1$?

- A. $y(t) = 2e^t - 1$.
 - B. $y(t) = 2e^{t+1} - 1$.
 - C. $y(t) = 2e^{t^2} - 1$.
 - D. $y(t) = \frac{1}{t+1}$.
 - E. $y(t) = \frac{2}{t+1}$.
 - F. $y(t) = \frac{1}{(t+1)^2}$.
10. (6 points) Which of the following computations are possible if A is an $(n \times m)$ matrix, B an $(n \times m)$ matrix, and C an $(n \times r)$ matrix, with $n \neq m \neq r$ (positive) integers.
- A. AB .
 - B. AB^TC .
 - C. AC^T .
 - D. ACB .
 - E. CA .
 - F. A^TC .

11. (4 points) Find the inverse of $A = \begin{bmatrix} 0 & 1 & -1 \\ 1 & 0 & 1 \\ 2 & 2 & 1 \end{bmatrix}$.

A. $A^{-1} = \begin{bmatrix} -2 & -3 & 2 \\ -1 & -2 & -1 \\ 2 & -2 & -2 \end{bmatrix}$

B. $A^{-1} = \begin{bmatrix} 2 & 3 & -1 \\ -1 & -2 & 1 \\ -2 & -2 & 1 \end{bmatrix}$

C. $A^{-1} = \begin{bmatrix} -2 & -3 & 1 \\ 1 & -2 & -1 \\ 2 & -2 & 2 \end{bmatrix}$

D. $A^{-1} = \begin{bmatrix} 2 & 3 & -1 \\ 1 & -2 & -2 \\ -2 & -2 & 1 \end{bmatrix}$

E. $A^{-1} = \begin{bmatrix} -2 & -3 & 1 \\ -1 & -2 & 1 \\ 2 & -2 & 1 \end{bmatrix}$

F. $A^{-1} = \begin{bmatrix} 2 & 3 & 1 \\ 1 & 2 & 1 \\ 2 & 2 & 1 \end{bmatrix}$

12. (4 points) What is the cross product between $u = (2, -3, 1)^T$ and $v = (1, 4, 5)^T$?

A. $u \times v = (19, 9, 10)$

B. $u \times v = (-9, -19, 11)$

C. $u \times v = (-19, -9, 11)$

D. $u \times v = (11, 19, -9)$

E. $u \times v = (10, 9, -19)$


F. None of the above.

13. (25 points)

We consider the function

$$f(x) = \frac{3x^2 - 1}{x^2 - 1}.$$

- 2 (a) What is the domain of the function?
- 3 (b) What are the intercepts with the x -axis and with the y -axis?
- 2 (c) What are the horizontal asymptotes?
- 2 (d) What are the vertical asymptotes?
- 6 (e) Compute and analyze the first derivative. In which intervals is the function increasing or decreasing, what are the local minima or maxima?
- 5 (f) Compute and analyze the second derivative. In which intervals is the function concave up or concave down, what are the points of inflection?
- 5 (g) Sketch the function. Your drawing needs to include all the qualitative features of the graph discussed in the questions above.



14. (25 points)

A specific plane in \mathbb{R}^3 is spanned by the three points $P = (2, 0, 1)$, $Q = (2, -2, 1)$ and $R = (3, 3, -1)$.

- 7 (a) Find two linearly independent vectors that are parallel to the plane. Briefly explain why your two vectors are linearly independent from each other.
- 9 (b) What method can you use to find a third vector that is orthogonal to two such linearly independent vectors? Using the vectors in (b), find a vector that is orthogonal to the plane.
- 9 (c) Write down the equation of the plane in normal form.